

Transmissible Spongiform Encephalopathies of Animals

Last Updated: October 2016

Importance

Transmissible spongiform encephalopathies (TSEs) are neurodegenerative diseases caused by prions. Although these infections usually remain asymptomatic for years, the disease is always progressive and fatal once the clinical signs develop. TSEs affecting animals include scrapie (tremblante de mouton, rida), bovine spongiform encephalopathy (BSE, “mad cow disease”), feline spongiform encephalopathy (FSE), transmissible mink encephalopathy (TME, mink scrapie) and chronic wasting disease (CWD). Although some prion diseases usually occur in one or a few closely related species, other prions can cross species barriers. BSE has a particularly wide host range. Cattle are the primary hosts for this disease, but some other ruminants, cats, lemurs and humans can also be affected; in cats, the disease is known as feline spongiform encephalopathy and in humans, it is called variant Creutzfeldt-Jakob disease (vCJD). Some evidence also suggests that TME might be caused by an atypical variant of the BSE agent called L-BSE. The discovery that BSE can cross species barriers and is zoonotic has raised concerns about all TSEs. Prion diseases in livestock can also result in trade sanctions. For these reasons, many countries are conducting control or eradication programs.

This document contains general information about the various prion diseases found in animals. Additional details can be found in the individual factsheet for each disease.

Etiology

Transmissible spongiform encephalopathies are caused by prions, infectious proteins that appear to replicate by converting a normal cellular protein into copies of the prion. The cellular protein, which is called PrP^c, is found on the surface of neurons. Pathogenic isoforms of PrP^c are designated PrP^{res} (‘res’ refers to the proteinase K-resistant nature of prions, compared to normal PrP^c). Alternative names for this protein are PrP^{TSE}, PrP^{Sc} (‘Sc’ stands for scrapie, the first animal prion disease to be recognized) and PrP^d (for disease-associated). Different prions (or, more accurately, different strains of PrP^{res}) cause scrapie, bovine spongiform encephalopathy (BSE) and chronic wasting disease (CWD). BSE in felids is called feline spongiform encephalopathy (FSE), and BSE in zoo ruminants was formerly called ‘spongiform encephalopathy of exotic ruminants.’

There are both classical and atypical BSE and scrapie prions. Atypical prions were first recognized during the last 20 years, as the result of enhanced surveillance for TSEs. The term ‘classical’ is used to distinguish the usual BSE and scrapie strains from these newly recognized forms. There are at least two atypical BSE prions. One has higher molecular mass fragments than classical BSE in an immunoblot and is called H-type BSE or H-BSE; the other has a lower molecular mass and is called L-type BSE or L-BSE. Currently, the most likely hypothesis is that L-BSE and H-BSE arise spontaneously in cattle, as genetic diseases. One of them may have given rise to the classical BSE epidemic after amplification through the food chain. Atypical/ Nor98 scrapie might, similarly, be a spontaneously occurring prion disease of small ruminants; however, this is not yet certain.

Which prion causes transmissible mink encephalopathy (TME) is still uncertain; however, it is expected to be an agent that occurs in meat fed to ranched mink. Current evidence suggests that it may be atypical L-BSE, although other possibilities, such as certain strains of the scrapie prion, have not been entirely ruled out.

Species Affected

Most prions seem to occur naturally in a small number of related species. Scrapie can affect sheep, goats, mouflon (*Ovis musimon*), and possibly other animals closely related to sheep and goats. Atypical/Nor98 scrapie has been found in both sheep and goats. CWD appears to be limited to cervids. As of 2016, it has been reported in free-living and/or captive mule deer (*Odocoileus hemionus*), black-tailed deer (*O. hemionus columbianus*), white-tailed deer (*O. virginianus*), Rocky Mountain elk (*Cervus elaphus nelsoni*), red deer (*C. elaphus elaphus*), sika deer (*C. nippon*), moose (*Alces alces*) and reindeer (*Rangifer tarandus tarandus*). TME has been seen only in ranched mink (*Neovision (Mustela) vision*). In contrast, the host range of the classical BSE prion



IOWA STATE UNIVERSITY
College of Veterinary Medicine



Transmissible Spongiform Encephalopathies

seems to be unusually broad. While cattle are the primary hosts, this prion has been found in housecats and captive zoo felids (where it is known as FSE), various species of exotic ruminants in zoos, two lemurs in a French zoo, and at least two goats. H-BSE and L-BSE have only been detected in cattle, to date, although some researchers suspect that L-BSE might cause TME in mink.

Some prions have been transmitted experimentally to additional species by routes that could occur in nature. Animals infected by feeding BSE prions include sheep, mink, cynomolgus macaques (*Macaca fascicularis*) and European red deer, but not pigs. Red deer did not appear to be very susceptible: only one of 6 animals became infected after they were fed a high dose of prions. Lemurs could be infected with L-BSE by the oral route, but little is otherwise known about animals' susceptibility to H-BSE or L-BSE. Scrapie does not seem to be transmitted readily to animals other than small ruminants. Squirrel monkeys (*Saimiri sciureus*) became infected when they were fed tissues that contained hamster-adapted scrapie prions, but several other species of primates, cattle and pigs did not appear to be susceptible. One fish (sea bream, *Sparus aurata*) had evidence of infection after oral inoculation with either BSE or scrapie prions. CWD can infect additional cervids and squirrel monkeys by the oral route, but attempts to infect cattle, cats, ferrets, mink and cynomolgus macaques have failed. Epidemiological studies also suggest that cattle are unlikely to be susceptible to this agent. Studies to date have found no evidence of CWD in wild predators and scavengers, such as coyotes (*Canis latrans*), mink, opossums (*Didelphis virginiana*) and raccoons (*Procyon lotor*), in endemic areas. There are few published experiments with TME; however, raccoons can be infected by either oral or parenteral routes.

Most prions have been transmitted to various animals other than their natural hosts by intracerebral inoculation. Because this route bypasses normal species barriers, it does not necessarily indicate that a species would be susceptible to that prion in nature. However, the absence of replication suggests it is unlikely to be susceptible. For instance, cats were resistant to scrapie via intracerebral inoculation, and CWD could not be transmitted to cynomolgus macaques or raccoons.

Zoonotic potential

Humans occasionally develop variant Creutzfeldt-Jakob disease (vCJD) after eating prion-containing tissues from an infected animal. To date, all known cases have been caused by the classical BSE prion, and the zoonotic potential of H-BSE and L-BSE is unclear. While FSE is also caused by the classical BSE prion, no cases of vCJD appear to have been acquired from cats. There is one curious case, where spongiform encephalopathies were reported simultaneously in a cat and its owner in 1998, and the prions found in both man and cat appeared to be similar. However, these prions differed from BSE, and the man was subsequently found to have the sporadic form of Creutzfeldt-Jakob disease, which is thought to be a spontaneous genetic disease.

As of 2016, investigations of suspicious cases of neurological disease in humans, surveillance, and epidemiological studies have found no evidence that CWD can affect humans. Nevertheless, the possibility that this agent is zoonotic cannot be ruled out at this time. People are not thought to be susceptible to scrapie, a disease that has been known since the 18th century.

Geographic Distribution

Animal prion diseases resemble neurological diseases caused by other agents, and information about their distribution can be incomplete outside regions with good veterinary surveillance. Classical scrapie has been reported on all major continents and some islands, although some countries recently found few or no cases during active surveillance. Australia and New Zealand have remained scrapie-free, with the exception of an outbreak caused by imported animals in the 1950s. Cases of classical BSE have been reported in indigenous cattle in some European countries, Canada, Israel and Japan. Some of these countries may have eradicated BSE, as it has not been detected in some time. Some other nations, including the U.S., have found this agent only in imported cattle. FSE has been reported in some countries where BSE occurs, and in housecats or zoo cats imported from these countries. Most infected animals had lived in the U.K. during the BSE epidemic.

Atypical BSE prions and atypical/ Nor98 scrapie prions have been reported in Europe, North America (both Canada and the U.S.) and some other countries, as the result of surveillance programs for TSEs. Both of these prions have been found in countries that are free of the classical form of that disease. Their presence does not affect a nation's scrapie or BSE status for international trade.

CWD has been found mainly in North America. It has been spreading from its original, limited focus in the U.S., and now occurs in a number of U.S. states and parts of Canada. This disease was imported to the Republic of Korea in captive cervids in 2001, and has been found there in captive animals, most recently in 2010. In 2016, CWD was detected in wild cervids in Norway. TME has been reported in ranched mink in the U.S., Canada, some European countries and the former U.S.S.R.

Transmission

TSEs are thought to be acquired primarily by ingestion, although some prions may also enter the body by other routes. Prions occur mainly in the central nervous system (CNS) of TSE-infected animals, but they can also be found in lymphoid tissues, peripheral nerves and various organs, to a greater or lesser extent depending on the prion and host species. Animals carry prions for life, and can transmit them even if they remain asymptomatic. Prions have also been transmitted iatrogenically in some species (e.g., via contaminated surgical instruments or in blood transfusions).

BSE, FSE and TME do not seem to be transmitted horizontally between animals. Animals (including cats with FSE) become infected with classical BSE prions when they

Transmissible Spongiform Encephalopathies

eat contaminated tissues from infected animals. BSE prions caused epidemics in cattle when they were amplified by recycling tissues from infected cattle into ruminant feed supplements. Young animals appear to be infected more readily than cattle over the age of 6 months. Vertical transmission has been reported in experimentally infected sheep, but it has not been documented in cattle. However, the offspring of BSE-infected cows seem to have an elevated risk of developing BSE, for reasons that are still unclear. There is one report of possible maternal transmission of FSE in a cheetah. TME is, likewise, thought to spread primarily by ingestion, probably when mink are fed tissues from infected animals; however, some evidence suggests that this prion might also enter the body through breaks in the mucous membranes or skin. Vertical transmission does not seem to occur in mink.

Scrapie and CWD can be transmitted directly between animals, or indirectly via the environment. CWD prions have been found, though sometimes at very low levels, in the saliva, blood, urine, feces and antler velvet of cervids. Most animals are thought to acquire this disease from the environment, probably during grazing. Vertical transmission has been documented in some cervid species. Most small ruminants seem to acquire scrapie from their dam during the neonatal period, although they can also become infected from unrelated animals or the environment. Older animals remain susceptible, though to a lesser extent. The placenta can contain scrapie prions, with particularly large amounts in sheep, and newborns may be infected by licking fetal membranes and fluids. The milk and colostrum of small ruminants are also infectious. *In utero* infection appears possible in sheep, though rare. Highly sensitive techniques have found low levels of scrapie prions in the urine, saliva and feces of sheep, but how much these sources contribute to transmission is still uncertain. The epidemiology of atypical/Nor98 scrapie suggests that it is not transmitted efficiently (or perhaps at all) between animals in nature, although laboratory experiments have demonstrated that newborn lambs are susceptible to oral inoculation.

Prions can persist for long periods in the environment. Infectious scrapie and CWD prions have been found in soil for 1.5 to 3 years or more in laboratory experiments. Evidence from Iceland, where scrapie-affected premises are not restocked for at least 2 years after depopulation and decontamination, suggests that this prion can infect animals up to 16 years later, under some conditions. Soil-bound prions remain infectious for animals. Repeated cycles of wetting and drying are reported to decrease, though not necessarily eliminate, soil infectivity. Prions can also remain infectious after passage through the digestive tracts of mammals and birds that eat infected animals.

Zoonotic transmission

In humans, variant Creutzfeldt-Jakob disease usually results from the ingestion of BSE prions, but a few people have been infected in blood transfusions from asymptomatic individuals. Other potential routes of infection include

organ/tissue transplantation and surgical transmission via contaminated equipment. Prions do not spread between people during casual contact.

Disinfection

Complete decontamination of prion-contaminated tissues, surfaces and environments can be difficult. These agents are very resistant to most disinfectants, including formalin and alcohol. They are also resistant to heat, or ultraviolet, microwave and ionizing radiation, particularly when they are protected in organic material or preserved with aldehyde fixatives, or when the prion titer is high. Prions can bind tightly to some surfaces, including stainless steel and plastic, without losing infectivity. Prions bound to metal seem to be highly resistant to decontamination.

Relatively few prion decontamination techniques have been published and confirmed to be effective for routine use. Some laboratories pre-treat tissues with formic acid to decrease infectivity before sectioning tissue blocks. A 1-2 N sodium hydroxide solution, or a sodium hypochlorite solution containing at least 2% (20,000 ppm) available chlorine, has traditionally been recommended for equipment and surfaces. Surfaces should be treated for more than 1 hour at 20°C (68°F). Overnight disinfection is recommended for equipment. Cleaning before disinfection removes organic material that may protect prions. Experimentally, some milder treatments have also been effective against certain prions, under some conditions. They include a specific phenolic disinfectant, various alkaline and enzymatic detergents (although the efficacy of specific agents within these classes varies), hydrogen peroxide gas plasma, radiofrequency gas plasma, and sodium dodecyl sulfate plus acetic acid. These agents might be useful for items that cannot withstand harsher decontamination procedures.

Physical inactivation of prions can be carried out by porous load autoclaving at 134°C (273°F) for 18 minutes at 30 lb/in². Resistance to heat may vary with the specific prion, the degree of contamination and type of sample. Tissue films containing prions are more difficult to decontaminate by steam after they have dried, and human guidelines for surgical instruments recommend that, after use, they be kept moist or wet until decontamination is performed. The cleaning agent used before autoclaving should also be chosen with care, as certain agents (e.g., some enzymatic treatments) can increase the resistance of prions to steam sterilization. Dry heat is less effective than moist heat; some prions can survive dry heat at temperatures as high as 360°C (680°F) for an hour, and one group even reported that infectivity survived incineration at 600°C (1112°F). A combination of chemical and physical decontamination can be more effective than either procedure alone, and effective combinations of chemical agents (e.g., NaOH) and autoclaving have been published. It should be noted that even the harshest combination of chemical and physical disinfection is not guaranteed to destroy all prions in all types of samples.

Transmissible Spongiform Encephalopathies

Decontaminating contaminated facilities, especially sites such as animal pens, may be very difficult. In one study, genetically susceptible sheep became infected with scrapie after being placed in pens that had been pressure washed and decontaminated with high concentrations of sodium hypochlorite (20,000 ppm free chlorine solution) for one hour, followed by painting and full re-galvanization or replacement of metalwork. Decontaminating soil contaminated with prions is currently impractical, although some agents, including an aqueous subtilisin-based enzymatic treatment (effective at ambient temperatures), appear promising in the laboratory. Incineration is commonly used for carcasses, but two studies found that composting may reduce or eliminate scrapie and other prions in tissues, while another suggested that soil microorganisms might degrade prions in buried carcasses.

Infections in Animals

Incubation Period

Transmissible spongiform encephalopathies have incubation periods of months or years. TME has the shortest estimated incubation period, at 6-12 months, while scrapie, BSE, FSE and CWD usually become apparent after 2 years or more in their natural hosts. Clinical cases can occur before this time, but are uncommon.

Clinical Signs

Transmissible spongiform encephalopathies are usually insidious in onset and tend to progress slowly. In most of these diseases, the clinical signs primarily involve the nervous system, and often include changes in behavior, as well as frank neurological signs such as tremors, ataxia and hyperreactivity to stimuli. Intense pruritus is common in sheep with classical scrapie; however, it tends to be less severe or absent in goats with this disease, and it is minimal or uncommon in most sheep or goats infected with atypical scrapie prions. Pruritus is not a common sign in BSE, CWD or TME. Loss of condition can be seen with all TSEs; however, wasting is particularly prominent in cervids with CWD, and some animals become severely emaciated before they die. Some cervids present mainly with weight loss, and few or subtle neurological signs. Once an animal becomes symptomatic, all TSEs are relentlessly progressive and fatal, typically within weeks to months.

Post Mortem Lesions [Click to view images](#)

There are no pathognomonic gross lesions in TSE-infected animals, although non-specific lesions may be seen. Animals that die or are euthanized in the early stages of these diseases may be in good condition; however, loss of condition is common later, and wasting of the carcass is often prominent in cervids with CWD. CWD can also result in megaesophagus and aspiration bronchopneumonia (which may be the cause of death), and classical scrapie-infected animals may have evidence of pruritus.

The typical histopathological lesions of TSEs are confined to the CNS. These lesions are usually (though not always) bilaterally symmetrical, and are characterized by non-inflammatory spongiform changes, with neuronal vacuolation and varying degrees of astrogliosis. Amyloid plaques may be seen in L-BSE and some cases of scrapie or CWD, but they are not usually found in other prion diseases, including classical BSE and H-BSE.

Diagnostic Tests

All TSEs can be diagnosed by examining the brain for prions at necropsy. In most diseases, prions can be detected in the medulla oblongata at the level of the obex; however, atypical/Nor98 scrapie can be missed by sampling only this site, and is more likely to be found in the cerebellum and other CNS locations. Asymptomatic ruminants can be tested for BSE or scrapie at routine slaughter by sampling the brainstem through the foramen magnum. TSEs may also be detected in lymphoid tissues at necropsy; in particular, scrapie and CWD can sometimes be found in these tissues before they are detected in the brain, and before the clinical signs appear.

Classical scrapie and CWD can also be diagnosed in live animals with biopsies of lymphoid tissues. Standardized sampling sites include the rectoanal mucosa-associated lymphoid tissue and palatine tonsil in both diseases, and the nictitating membrane (third eyelid) in scrapie. Rectal mucosa and nictitating membrane biopsies can be taken without sedation, using only topical anesthesia and restraint, and are more practical than tonsil biopsies in the field. Prions may occasionally be found in other lymphoid tissues, such as superficial lymph nodes. Live animal tests do not appear to be capable of detecting animals with atypical/Nor98 scrapie.

Immunoblotting (Western blotting) or immunohistochemistry are the most specific assays for detecting prions, and can be used in all of the animal prion diseases. A number of rapid diagnostic tests based on enzyme-linked immunosorbent assays (ELISAs), automated immunoblotting or other techniques (e.g., lateral flow assays) have also been validated for BSE and classical scrapie. Some of these tests appear to be capable of detecting other TSEs, though with varying efficacy. Rapid tests allow large numbers of samples to be screened, and are often used in surveillance and slaughter testing. Histological examination of the brain can be very helpful in the diagnosis of prion diseases (although it is not generally used as the sole confirmatory test), but some animals in the early stages of infection have few or no spongiform changes. The presence of characteristic prion fibrils in the brain, called scrapie-associated fibrils (SAF), may be used to diagnose TSEs with electron microscopy, even in autolyzed brains; however, this test has low sensitivity.

Highly sensitive assays, including protein misfolding cyclic amplification (PMCA) and quaking-induced conversion (QuIC) or real-time quaking-induced conversion (RT-QuIC), may be able to identify infected animals earlier than immunoblotting or immunohistochemistry. They can also find prions in tissues where they are present in very low

Transmissible Spongiform Encephalopathies

amounts. These techniques detect tiny amounts of prions by their ability to convert PrP^c (the normal cellular protein) into prions *in vitro*. They are mainly used in research at present, but are being investigated for possible diagnostic use in several diseases. TSEs can also be detected by inoculation into mice (rodent bioassays); however, an incubation period of several months makes this technique impractical for routine diagnosis. Serology is not useful for diagnosis, as antibodies are not made against prions.

Treatment

There is no treatment for any prion disease.

Control

Disease reporting

Veterinarians who encounter or suspect a prion disease should follow their national and/or local guidelines for disease reporting. Diseases that are absent from a nation are reportable immediately in most countries. Endemic diseases can also be reportable, although the requirements may be different.

In the U.S., state or federal veterinary authorities should be informed immediately of any cases of BSE, FSE or TME. The U.S. has an eradication program for scrapie, and this disease is also reportable. Requirements for CWD may vary with the state, and with a farmed cervid herd's participation in the U.S. herd certification program. Local wildlife management agencies are often responsible for CWD surveillance programs in wild cervids.

Prevention

BSE, FSE and TME can be prevented by not feeding tissues that may contain prions to susceptible species. Complete avoidance is necessary, as cooking or rendering cannot completely inactivate these agents. Banning ruminant (or mammalian) tissues from ruminant (or animal) feed has significantly reduced the number of new cases of BSE and controlled the epidemics in cattle. The specific bans, and protein sources prohibited, vary with the country. Surveillance and tracing of infected animals can reveal cohorts that may have received the same feed. Countries may place trade bans on the importation of live animals and certain animal proteins from TSE-affected nations.

Diseases that can spread horizontally, such as scrapie and CWD, require additional methods of control. The risk of introducing these diseases can be reduced by maintaining a closed herd/ flock or minimizing outside purchases of stock. If replacement animals must be added, they should be from herds that test negative for the disease and are managed in a way that makes them unlikely to become infected. Milk and colostrum from potentially infected sheep or goats should not be fed to scrapie-free flocks. Reducing exposure to high concentrations of prions (e.g., in the placenta of sheep) may reduce transmission within a herd or flock that has become infected. The risk that a sheep will develop scrapie is strongly influenced by its genotype, and selecting genetically resistant animals can aid control efforts. Whether genetic control

methods are feasible in scrapie-infected goat herds or CWD-infected cervid herds is still uncertain. Complete depopulation, followed by cleaning and disinfection, is sometimes used on prion-infected farms; however, decontamination may be difficult and the disease may recur. Some countries have developed voluntary and/or mandatory programs to control CWD and scrapie, with eradication from domestic herds/ flocks as the ultimate goal.

Controlling CWD in wild cervids is very difficult. Infected captive cervids should be kept from contact with wild cervids, to avoid transmitting this disease to uninfected areas. Many states and provinces also have restrictions on transporting tissues from hunter-killed cervids in CWD-endemic areas. Attempts to reduce transmission in wild populations include banning practices that encourage cervids to congregate in certain areas (e.g., feeding or baiting), and culling to reduce animal densities. Other TSEs are not known to be an issue in wild animals.

There are no established control methods for L-BSE, H-BSE or atypical/ Nor98 scrapie, which could be spontaneous genetic diseases. These prions do not currently affect a nation's BSE or scrapie status for international trade.

Morbidity and Mortality

TSEs are usually diseases of adult animals. Only adult mink are usually affected by TME; in at least one outbreak, kits housed with their dams and eating the same diet were completely spared. The peak incidence of classical BSE occurs in 4-6 year-old cattle, while classical scrapie peaks in sheep between the ages of 2 and 5 years, and CWD in captive cervids between 2 and 7 years. All of the ruminant and cervid TSEs are very rare or absent in animals that are less than 12-18 months of age. L-BSE, H-BSE and atypical/ Nor98 scrapie tend to occur in animals older than those with classical BSE or classical scrapie, respectively. All TSEs are fatal once clinical signs appear.

The animal's genotype influences the onset and severity of some TSEs. This is particularly well-known in sheep, where susceptibility or resistance to classical scrapie is associated with polymorphisms in the PrP gene at codons 136, 154 and 171. Sheep with resistant genotypes become ill after longer incubation periods, or not at all. A genetically resistant fetus also suppresses the appearance of prions in the placenta of its genetically-susceptible, infected dam. The genes that affect scrapie resistance are not as well understood in goats, but some polymorphisms have been identified, and one (K222) has been proposed as a possible target for breeding more scrapie-resistant animals. Atypical scrapie is also affected by the animal's genotype; however, this disease seems to be common in sheep resistant to classical scrapie, and underreported in the animals most susceptible to the latter form. Genetic susceptibility to CWD is not yet well understood.

TSEs vary greatly in prevalence, depending on the country, host species and specific disease. Atypical BSE and atypical scrapie seem to arise sporadically, at low levels, in their respective host species. Atypical scrapie prions are typically found in only a single animal per flock or herd, while

Transmissible Spongiform Encephalopathies

only about 100 animals infected with L-BSE or H-BSE have been identified since surveillance began. TME also seems to be rare, with few reported outbreaks. However, large numbers of mink are often affected on infected ranches. FSE was documented in approximately a hundred housecats and twenty zoo felids between 1990 and 2007. Most of these cases occurred in the U.K. during the BSE epidemic, and new cases have not been published since 2007.

Classical BSE, classical scrapie and CWD can be common in some areas. Most BSE cases occurred during epidemics that began in the 1980s in the U.K., and resulted from recycling tissues from infected animals in ruminant feed. At one time, the estimated prevalence of classical BSE in infected countries ranged from more than 100 cases per million cattle to fewer than 2 cases per million. As a result of control measures (particularly feed bans), the incidence of this disease has declined worldwide. In the U.K., the number of new cases has dropped from nearly 1,000 per week, at the height of the epidemic in 1992, to 0-3 cases per year between 2012 and 2016.

Classical scrapie can be a significant problem in some areas, while other regions report few or no cases. This disease is much less common in goats than sheep. Between 2002 and 2009, surveillance programs in the E.U. identified approximately 3300 infected goats, compared to about 15,000 sheep. The average prevalence of scrapie in sheep was 0.087% in E.U. countries reporting this disease between 2002 and 2012, and 0.015% in the U.S., as of 2013. In North America, the prevalence of CWD in wild cervids is reported to be < 5% in deer and < 2.5% in elk in many affected areas. However, it differs between regions, and can be higher in localized "hot spots," where it may be present in up to 10-12% of elk and up to 50% of wild deer. Only a few CWD cases have been reported, to date, in wild cervids in Norway, and its prevalence there is uncertain.

Infections in Humans

BSE

BSE is the only animal TSE known to affect humans; people who have eaten BSE prions may develop variant Creutzfeldt-Jakob disease. As of May 2016, 228 cases of vCJD have been reported worldwide. With the exception of 27 cases in France, most of these people had resided in the U.K. for more than 6 months during the peak of the BSE epidemic, and are likely to have been infected there. In the U.K., the incidence of vCJD peaked in 2000, when 28 cases were diagnosed, and gradually fell to 2-5 cases per year between 2006 and 2011. Only two additional cases were recognized between 2012 and 2016. The number of people who have been infected asymptotically, and the percentage of those likely to develop vCJD, is still unclear. Based on the decreasing number of clinical cases, some sources suggest that few additional cases may be seen. However, some studies that have examined lymphoid tissues, such as the tonsils or appendix, suggest that from 1 in 2000 to 1 in 10,000 people in the U.K. may be infected

subclinically with BSE prions. Whether these people will ever develop vCJD is not known.

The symptoms of vCJD are broadly similar to the sporadic (genetic) form of Creutzfeldt-Jakob disease, but usually appear in younger patients. The median age of onset is 26 years, although cases have been reported in people from 12 to 74 years. The first signs are usually psychiatric symptoms such as anxiety, depression and social withdrawal, and/or persistent painful sensory symptoms. In most patients, frank neurological signs (e.g., ataxia, tremors) appear a few months later; however, neurological signs coincide with or precede psychiatric symptoms in a minority of patients. Cognitive function gradually deteriorates, and involuntary movements and visual disturbances may develop late in the course of the disease. There is no known treatment, and most patients die within 2 years. To date, all individuals with confirmed clinical cases have been homozygous for methionine at codon 129 in the PrP^C protein.

Prevention of vCJD is based on the avoidance of high-risk tissues from cattle. Many countries exclude animals with BSE-like clinical signs, as well as high risk tissues from asymptomatic cattle (specified risk materials or SRM), from human food. SRM generally include the brain, spinal cord, associated bones and some nerve ganglia; tonsils; and various portions of the intestinal tract (e.g., currently, the distal ileum in the U.S., and the last 4 meters of the small intestine, the cecum and mesentery in the E.U.). Some nations conduct active surveillance of cattle at slaughter, using rapid tests, to detect cases of BSE. Positive carcasses are destroyed. Slaughter and processing techniques that have a high risk of contaminating muscle tissues with CNS have also been prohibited in many nations. Many countries restrict blood donations from people with a significant risk of having been infected during the BSE epidemics. Some may also take other measures to prevent transmission in blood transfusions or during high-risk surgical procedures. Although laboratory or abattoir-related cases have not been reported, veterinarians and laboratory workers should always take precautions when conducting necropsies on BSE-suspects or handling tissues; BSL-3 is the recommended level of protection for handling these prions.

CWD

Although no human infections with CWD have ever been reported, the possibility that this disease could be zoonotic has not been ruled out. Hunters should consider having carcasses tested for CWD; information on this program is available from most state wildlife agencies in the U.S. Meat from cervids that appear to be ill, as well as meat from apparently healthy animals that test positive for CWD, should not be eaten or fed to any animals. Gloves should be worn when field-dressing a cervid carcass. Boning-out the meat and minimizing the handling of the brain, spinal cord and lymphoid tissues associated with the gastrointestinal tract (e.g. tonsils) from cervids may reduce the risk of exposure, but will not necessarily remove all prions.

Transmissible Spongiform Encephalopathies

Internet Resources

[Canadian Food Inspection Agency \(CFIA\). BSE](#)

[Canadian Food Inspection Agency. Chronic Wasting Disease \(CWD\) of Deer and Elk \(including information on the herd certification program\)](#)

[Centers for Disease Control and Prevention \(CDC\). Prion Diseases.](#)

[Chronic Wasting Disease Alliance](#)

[European Commission. Control of TSEs \(including BSE and scrapie\)](#)

[European Food Safety Authority. Transmissible Spongiform Encephalopathies](#)

[European Union Reference Laboratory, TSE-LAB-NET](#)

[Scrapie Canada](#)

[The National Creutzfeldt-Jakob Disease Surveillance Unit, United Kingdom.](#)

[United Kingdom. Department for Environment Food and Rural Affairs \(DEFRA\). Bovine Spongiform Encephalopathy](#)

[U.K. DEFRA. Exotic species and domestic cats: TSE surveillance statistics](#)

[United States Department of Agriculture \(USDA\). Animal and Plant Health Inspection Service \(APHIS\).](#)

[USDA APHIS. Chronic Wasting Disease \(including information on the CWD Certified and Monitored Herd Programs\).](#)

[USDA APHIS Scrapie Program](#)

[United States Food and Drug Administration \(FDA\). Bovine Spongiform Encephalopathy](#)

[U.S. Geological Survey \(USGS\). Chronic Wasting Disease](#)

[World Organization for Animal Health \(WOAH\)](#)

[WOAH Manual of Diagnostic Tests and Vaccines for Terrestrial Animals](#)

[WOAH Terrestrial Animal Health Code](#)

Acknowledgements

This factsheet was written by Anna Rovid Spickler, DVM, PhD, Veterinary Specialist from the Center for Food Security and Public Health. The U.S. Department of Agriculture Animal and Plant Health Inspection Service (USDA APHIS) provided funding for this factsheet through a series of cooperative agreements related to the development of resources for initial accreditation training.

The following format can be used to cite this factsheet. Spickler, Anna Rovid. 2016. *Transmissible Spongiform*

Encephalopathies. Retrieved from <http://www.cfsph.iastate.edu/DiseaseInfo/factsheets.php>.

References

- Acín C, Martín-Burriel I, Monleón E, Lyahyai J, Pitarch JL, Serrano C, Monzón M, Zaragoza P, Badiola JJ. Prion protein gene variability in Spanish goats. Inference through susceptibility to classical scrapie strains and pathogenic distribution of peripheral PrP(sc.). *PLoS One*. 2013;8(4):e61118.
- Acín C, Pitarch JL. Controlling scrapie and bovine spongiform encephalopathy in goats. *Vet Rec*. 2016;178(7):166-7.
- Acutis PL, Martucci F, D'Angelo A, Peletto S, Colussi S, et al. Resistance to classical scrapie in experimentally challenged goats carrying mutation K222 of the prion protein gene. *Vet Res*. 2012;43:8.
- Aguilar-Calvo P, Espinosa JC, Pintado B, Gutiérrez-Adán A, Alamillo E, Miranda A, Prieto I, Bossers A, Andreoletti O, Torres JM. Role of the goat K222-PrP(C) polymorphic variant in prion infection resistance. *J Virol*. 2014;88(5):2670-6.
- Aguzzi A, Heikenwalder M, Miele G. Progress and problems in the biology, diagnostics, and therapeutics of prion diseases. *J Clin Invest*. 2004;114:153-160.
- Alverson J, O'Rourke KI, Baszler TV. PrPSc accumulation in fetal cotyledons of scrapie-resistant lambs is influenced by fetus location in the uterus. *J Gen Virol*. 2006;87:1035-41.
- Anderson CA, Bosque P, Filley CM, Arciniegas DB, Kleinschmidt-Demasters BK, Pape WJ, Tyler KL. Colorado surveillance program for chronic wasting disease transmission to humans: lessons from 2 highly suspicious but negative cases. *Arch Neurol*. 2007;64:439-41.
- Andreoletti O, Lacroux C, Chabert A, Monnereau L, Tabouret G, Lantier F, Berthon P, Eychenne F, Lafond-Benestad S, Elsen JM, Schelcher F. PrP(Sc) accumulation in placentas of ewes exposed to natural scrapie: influence of foetal PrP genotype and effect on ewe-to-lamb transmission. *J Gen Virol*. 2002;83:2607-16.
- Andreoletti O, Morel N, Lacroux C, Rouillon V, Barc C, et al. Bovine spongiform encephalopathy agent in spleen from an ARR/ARR orally exposed sheep. *J Gen Virol*. 2006;87:1043-6.
- Andreoletti O, Orge L, Benestad SL, Beringue V, Litaize C, Simon S, Le Dur A, Laude H, Simmons H, Lugan S, Corbière F, Costes P, Morel N, Schelcher F, Lacroux C. Atypical/Nor98 scrapie infectivity in sheep peripheral tissues. *PLoS Pathog*. 2011;7(2):e1001285.
- Andreoletti O, Simon S, Lacroux C, Morel N, Tabouret G, Chabert A, Lugan S, Corbière F, Ferre P, Foucras G, Laude H, Eychenne F, Grassi J, Schelcher F. PrPSc accumulation in myocytes from sheep incubating natural scrapie. *Nat Med*. 2004;10:591-3.
- Andrews NJ. Incidence of variant Creutzfeldt-Jakob disease diagnoses and deaths in the UK January 1994 – December 2011 The National Creutzfeldt-Jakob Disease Surveillance Unit [CJD Unit], United Kingdom. Available at: <http://www.cjd.ed.ac.uk/data.html>. Accessed 28 Aug 2016.
- Angers RC, Browning SR, Seward TS, Sigurdson CJ, Miller MW, Hoover EA, Telling GC. Prions in skeletal muscles of deer with chronic wasting disease. *Science*. 2006;311:1117.

Transmissible Spongiform Encephalopathies

- Angers RC, Seward TS, Napier D, Green M, Hoover E, Spraker T, O'Rourke K, Balachandran A, Telling GC. Chronic wasting disease prions in elk antler velvet. *Emerg Infect Dis.* 2009;15(5):696-703.
- Animal Health Australia. TSE freedom assurance report. 2014-2015. Available at: <http://TSEFAP-2014-15-Final-Report-for-website.pdf>. Accessed 28 Sept 2016.
- Acquatella-Tran Van Ba I, Imberdis T, Perrier V. From prion diseases to prion-like propagation mechanisms of neurodegenerative diseases. *Int J Cell Biol.* 2013;2013:975832.
- Arnold ME, Wilesmith JW. Estimation of the age-dependent risk of infection to BSE of dairy cattle in Great Britain. *Prev Vet Med.* 2004;66:35-47.
- Arsac JN, Andreoletti O, Bilheude JM, Lacroux C, Benestad SL, Baron T. Similar biochemical signatures and prion protein genotypes in atypical scrapie and Nor98 cases, France and Norway. *Emerg Infect Dis.* 2007;13:58-65.
- Baeten LA, Powers BE, Jewell JE, Spraker TR, Miller MW. A natural case of chronic wasting disease in a free-ranging moose (*Alces alces shirasi*). *J Wildl Dis.* 2007;43:309-14.
- Balkema-Buschmann A, Eiden M, Hoffmann C, Kaatz M, Ziegler U, Keller M, Groschup MH. BSE infectivity in the absence of detectable PrP(Sc) accumulation in the tongue and nasal mucosa of terminally diseased cattle. *J Gen Virol.* 2011;92(Pt 2):467-76.
- Balkema-Buschmann A, Fast C, Kaatz M, Eiden M, Ziegler U, McIntyre L, Keller M, Hills B, Groschup MH. Pathogenesis of classical and atypical BSE in cattle. *Prev Vet Med.* 2011;102(2):112-7.
- Balkema-Buschmann A, Ziegler U, McIntyre L, Keller M, Hoffmann C, Rogers R, Hills B, Groschup MH. Experimental challenge of cattle with German atypical bovine spongiform encephalopathy (BSE) isolates. *J Toxicol Environ Health A.* 2011;74(2-4):103-9.
- Balter M. Intriguing clues to a scrapie-mad cow link. *Science.* 2001;292:827-9.
- Bannach O, Birkmann E, Reinartz E, Jaeger KE, Langeveld JP, Rohwer RG, Gregori L, Terry LA, Willbold D, Riesner D. Detection of prion protein particles in blood plasma of scrapie infected sheep. *PLoS One.* 2012;7(5):e36620.
- Barlow RM. Transmissible mink encephalopathy: pathogenesis and nature of the aetiological agent. *J Clin Pathol Suppl (R Coll Pathol).* 1972;6:102-9.
- Baron T, Belli P, Madec JY, Moutou F, Vitaud C, Savey M. Spongiform encephalopathy in an imported cheetah in France. *Vet Rec.* 1997;141:270-1.
- Baron T, Bencsik A, Biacabe AG, Morignat E, Bessen RA. Phenotypic similarity of transmissible mink encephalopathy in cattle and L-type bovine spongiform encephalopathy in a mouse model. *Emerg Infect Dis.* 2007;13:1887-94.
- Baron T, Biacabe AG, Arsac JN, Benestad S, Groschup MH. Atypical transmissible spongiform encephalopathies (TSEs) in ruminants. *Vaccine.* 2007;25:5625-30.
- Baron T, Vulin J, Biacabe AG, Lakhdar L, Verchere J, Torres JM, Bencsik A. Emergence of classical BSE strain properties during serial passages of H-BSE in wild-type mice. *PLoS One.* 2011 Jan 14;6(1):e15839.
- Barria MA, Ironside JW, Head MW. Exploring the zoonotic potential of animal prion diseases: *in vivo* and *in vitro* approaches. *Prion.* 2014;8(1):85-91.
- Bartz JC, Kincaid AE, Bessen RA. Rapid prion neuroinvasion following tongue infection. *J Virol.* 2003;77:583-91.
- Bartz JC, McKenzie DI, Bessen RA, Marsh RF, Aiken JM. Transmissible mink encephalopathy species barrier effect between ferret and mink: PrP gene and protein analysis. *J Gen Virol.* 1994;75:2947-53.
- Beekes M, McBride PA. The spread of prions through the body in naturally acquired transmissible spongiform encephalopathies. *FEBS J.* 2007;274:588-605.
- Belay ED, Blase J, Schulster LM, Maddox RA, Schonberger LB. Management of neurosurgical instruments and patients exposed to Creutzfeldt-Jakob disease. *Infect Control Hosp Epidemiol.* 2013;34(12):1272-80.
- Belay ED, Maddox RA, Williams ES, Miller MW, Gambetti P, Schonberger LB. Chronic wasting disease and potential transmission to humans. *Emerg Infect Dis.* 2004;10:977-84.
- Bellworthy SJ, Dexter G, Stack M, Chaplin M, Hawkins SA, Simmons MM, Jeffrey M, Martin S, Gonzalez L, Hill P. Natural transmission of BSE between sheep within an experimental flock. *Vet Rec.* 2005;157:206.
- Belondrade M, Nicot S, Béringue V, Coste J, Lehmann S, Bougard D. Rapid and highly sensitive detection of variant Creutzfeldt-Jakob disease abnormal prion protein on steel surfaces by protein misfolding cyclic amplification: application to prion decontamination studies. *PLoS One.* 2016;11(1):e0146833.
- Bencsik A, Debeer S, Petit T, Baron T. Possible case of maternal transmission of feline spongiform encephalopathy in a captive cheetah. *PLoS One.* 2009;4(9):e6929.
- Benestad SL, Arsac JN, Goldmann W, Nöremark M. Atypical/Nor98 scrapie: properties of the agent, genetics, and epidemiology. *Vet Res.* 2008;39(4):19.
- Benestad SL, Sarradin P, Thu B, Schonheit J, Tranulis MA, Bratberg B. Cases of scrapie with unusual features in Norway and designation of a new type, Nor98. *Vet Rec.* 2003;153:202-8.
- Béringue V, Andréoletti O, Le Dur A, Essalmani R, Vilotte JL, Lacroux C, Reine F, Herzog L, Biacabé AG, Baron T, Caramelli M, Casalone C, Laude H. A bovine prion acquires an epidemic bovine spongiform encephalopathy strain-like phenotype on interspecies transmission. *J Neurosci.* 2007;27:6965-71.
- Bessen RA, Marsh RF. Biochemical and physical properties of the prion protein from two strains of the transmissible mink encephalopathy agent. *J Virol.* 1992;66:2096-101.
- Bessen RA, Shearin H, Martinka S, Boharski R, Lowe D, Wilham JM, Caughey B, Wiley JA. Prion shedding from olfactory neurons into nasal secretions. *PLoS Pathog.* 2010;6:1000837.
- Boujon C, Serra F, Seuberlich T. Atypical variants of bovine spongiform encephalopathy: rare diseases with consequences for BSE surveillance and control. *Schweiz Arch Tierheilkd.* 2016;158(3):171-7.
- Bozzetta E, Nappi R, Crudeli S, Meloni D, Varello K, Loprevite D, Melis PG, Mazza M, Colussi S, Ingravalle F, Ru G, Nonno R, Ligios C. Comparative performance of three TSE rapid tests for surveillance in healthy sheep affected by scrapie. *Virol Methods.* 2011;173(2):161-8.

Transmissible Spongiform Encephalopathies

- Bradshaw JM, Pearson GR, Gruffydd-Jones TJ. A retrospective study of 286 cases of neurological disorders of the cat. *J Comp Pathol*. 2004;131:112-20.
- Brandt AL, Kelly AC, Green ML, Shelton P, Novakowski J, Mateus-Pinilla NE. Prion protein gene sequence and chronic wasting disease susceptibility in white-tailed deer (*Odocoileus virginianus*). *Prion*. 2015;9(6):449-62.
- Brown P, Abee CR. Working with transmissible spongiform encephalopathy agents. *ILAR J*. 2005;46:44-52.
- Brown P, Gajdusek DC. Survival of scrapie virus after 3 years' interment. *Lancet*. 1991;337:269-70.
- Brown P, McShane LM, Zanusso G, Detwile L. On the question of sporadic or atypical bovine spongiform encephalopathy and Creutzfeldt-Jakob disease. *Emerg Infect Dis*. 2006;12:1816-21.
- Brown P, Rau EH, Johnson BK, Bacote AE, Gibbs CJ, Jr., Gajdusek DC. New studies on the heat resistance of hamster-adapted scrapie agent: threshold survival after ashing at 600 degrees C suggests an inorganic template of replication. *Proc Natl Acad Sci USA*. 2000;97:3418-21.
- Bulgin MS. Overview of scrapie. Kahn CM, Line S, Aiello SE, editors. *The Merck veterinary manual* [online]. Whitehouse Station, NJ: Merck and Co; 2013. Scrapie. Available at: http://www.merckvetmanual.com/mvm/nervous_system/scrapie/overview_of_scrapie.html. Accessed 30 Aug 2016.
- Buschmann A, Biacabe AG, Ziegler U, Bencsik A, Madec JY, Erhardt G, Luhken G, Baron T, Groschup MH. Atypical scrapie cases in Germany and France are identified by discrepant reaction patterns in BSE rapid tests. *J Virol Methods*. 2004;117:27-36.
- Buschmann A, Luhken G, Schultz J, Erhardt G, Groschup MH. Neuronal accumulation of abnormal prion protein in sheep carrying a scrapie-resistant genotype (PrPARR/ARR). *J Gen Virol*. 2004;85:2727-33.
- Canadian Food Inspection Agency (CFIA). Enhanced animal health protection from BSE - specified risk material (SRM). CFIA; 2015 Apr. Available at: <http://www.inspection.gc.ca/animals/terrestrial-animals/diseases/reportable/bse/srm/eng/1299870250278/1334278201780>. Accessed 28 Aug 2016.
- Carmona P, Monzon M, Monleon E, Badiola JJ, Monreal J. *In vivo* detection of scrapie cases from blood by infrared spectroscopy. *J Gen Virol*. 2005;86:3425-31.
- Centers for Disease Control and Prevention (CDC). Chronic wasting disease [online]. CDC; 2007 Jan. Available at: <http://www.cdc.gov/ncidod/dvrd/cwd>. Accessed 19 Sept. 2008.
- Centers for Disease Control and Prevention [CDC]. Variant Creutzfeldt-Jakob disease [website online]. CDC; 2015 Feb. Available at: <http://www.cdc.gov/prions/vcjd/index.html>. Accessed 28 Aug 2016.
- Chang B, Cheng X, Yin S, Pan T, Zhang H, Wong P, Kang SC, Xiao F, Yan H, Li C, Wolfe LL, Miller MW, Wisniewski T, Greene MI, Sy MS. Test for detection of disease-associated prion aggregate in the blood of infected but asymptomatic animals. *Clin Vaccine Immunol*. 2007;14:36-43.
- Chronic Wasting Disease Alliance. Carcass transportation regulations in the United States and Canada [online]. Available at: <http://www.cwd-info.org/index.php/fuseaction/policy.main>. Accessed 19 Sept 2008.
- Chronic Wasting Disease Alliance. Chronic wasting disease (CWD) outbreaks and surveillance program in the Republic of Korea. 2011. Available at: <http://cwd-info.org/chronic-wasting-disease-cwd-outbreaks-and-surveillance-program-in-the-republic-of-korea/>. Accessed 28 July 2016.
- Chronic Wasting Disease Alliance. The first detection of chronic wasting disease (CWD) in Europe. 2016. Available at: <http://cwd-info.org/the-first-detection-of-chronic-wasting-disease-cwd-in-europe/>. Accessed 29 Jul 2016.
- Colorado Division of Wildlife. Chronic wasting disease in Colorado, 2006–2008. Colorado Division of Wildlife Report. Fort Collins, Colorado; 2009. 5 p.
- Corbière F, Perrin-Chauvineau C, Lacroux C, Costes P, Thomas M, Brémaud I, Martin S, Lugan S, Chartier C, Schelcher F, Barillet F, Andreoletti O. PrP-associated resistance to scrapie in five highly infected goat herds. *Gen Virol*. 2013;94(Pt 1):241-5.
- Cunningham AA, Kirkwood JK, Dawson M, Spencer YI, Green RB, Wells GA. Bovine spongiform encephalopathy infectivity in greater kudu (*Tragelaphus strepsiceros*). *Emerg Infect Dis*. 2004;10:1044-9.
- Curcio L, Sebastiani C, Di Lorenzo P, Lasagna E, Biagetti M. Review: A review on classical and atypical scrapie in caprine: Prion protein gene polymorphisms and their role in the disease. *Animal*. 2016 Apr 25;1-9. [Epub ahead of print]
- Dagleish MP, Martin S, Steele P, Finlayson J, Eaton SL, Sisó S, et al. Susceptibility of European red deer (*Cervus elaphus elaphus*) to alimentary challenge with bovine spongiform encephalopathy. *PLoS One*. 2015;10(1):e0116094.
- Dassanayake RP, Orrú CD, Hughson AG, Caughey B, Graça T, Zhuang D, Madsen-Bouterse SA, Knowles DP, Schneider DA. Sensitive and specific detection of classical scrapie prions in the brains of goats by real-time quaking-induced conversion. *J Gen Virol*. 2016;97(3):803-12.
- Dassanayake RP, Truscott TC, Zhuang D, Schneider DA, Madsen-Bouterse SA, Young AJ, Stanton JB, Davis WC, O'Rourke KI. Classical natural ovine scrapie prions detected in practical volumes of blood by lamb and transgenic mouse bioassays. *J Vet Sci*. 2015;16(2):179-86.
- Daus ML, Breyer J, Wagenfuehr K, Wemheuer WM, Thomzig A, Schulz-Schaeffer WJ, Beekes M. Presence and seeding activity of pathological prion protein (PrP(TSE)) in skeletal muscles of white-tailed deer infected with chronic wasting disease. *PLoS One*. 2011;6(4):e18345.
- Dexter G, Tongue SC, Heasman L, Bellworthy SJ, Davis A, Moore SJ, Simmons MM, Sayers AR, Simmons HA, Matthews D. The evaluation of exposure risks for natural transmission of scrapie within an infected flock. *BMC Vet Res*. 2009;5:38.
- Diack AB, Head MW, McCutcheon S, Boyle A, Knight R, Ironside JW, Manson JC, Will RG. Variant CJD. 18 years of research and surveillance. *Prion*. 2014;8(4):286-95.
- Doherr MG. Brief review on the epidemiology of transmissible spongiform encephalopathies (TSE). *Vaccine*. 2007;25:5619-24.
- Eckroade RJ. 1972. Neuropathology and experimental transmission to other species of transmissible mink encephalopathy. PhD Thesis, University of Wisconsin, Madison, WI.

Transmissible Spongiform Encephalopathies

- Eckroade RJ, Zu Rhein GM, Hanson RP: Transmissible mink encephalopathy in carnivores: clinical, light and electron microscopic studies in raccoons, skunks and ferrets. *J Wildl Dis.* 1973;9:229-40.
- Eckroade RJ, Zu Rhein GM, Marsh RF, Hanson RP. Transmissible mink encephalopathy: experimental transmission to the squirrel monkey *Science.* 1970;169:1088-90.
- Edgeworth JA, Sicilia A, Linehan J, Brandner S, Jackson GS, Collinge J. A standardized comparison of commercially available prion decontamination reagents using the standard steel-binding assay. *J Gen Virol.* 2011;92(Pt 3):718-26.
- Eghiaian F, Grosclaude J, Lesceu S, Debey P, Doublet B, Tréguer E, Rezaei H, Knossow M. Insight into the PrPC -> PrPSc conversion from the structures of antibody-bound ovine prion scrapie-susceptibility variants *Proc Natl Acad Sci U S A.* 2004;101:10254-9.
- Eiden M, Hoffmann C, Balkema-Buschmann A, Müller M, Baumgartner K, Groschup MH. Biochemical and immunohistochemical characterization of feline spongiform encephalopathy in a German captive cheetah. *J Gen Virol.* 2010;91(Pt 11):2874-83.
- Eloit M, Adjou K, Coulpier M, Fontaine JJ, Hamel R, et al. BSE agent signatures in a goat. *Vet Rec.* 2005;156:523-4.
- Espinosa JC, Andréoletti O, Castilla J, Herva ME, Morales M, Alamillo E, San-Segundo FD, Lacroux C, Lugan S, Salguero FJ, Langeveld J, Torres JM. Sheep-passaged bovine spongiform encephalopathy agent exhibits altered pathobiological properties in bovine-PrP transgenic mice. *Virol.* 2007;81:835-43.
- Espinosa JC, Morales M, Castilla J, Rogers M, Torres JM. Progression of prion infectivity in asymptomatic cattle after oral bovine spongiform encephalopathy challenge. *J Gen Virol.* 2007;88:1379-83.
- European Commission. Control of TSEs (including BSE and scrapie). 2016. Available at: http://ec.europa.eu/food/safety/biosafety/food_borne_diseases/tse_bse/index_en.htm#. Accessed 26 Aug 2016.
- European Food Safety Authority [EFSA]. EFSA opinion on the likelihood of BSE infectivity in specified risk material. EFSA; 2007 Jul. Available at: http://www.efsa.europa.eu/en/press_room/press_release/pr_sr_m_back_calculation.html. Accessed 25 Aug 2007.
- European Food Safety Authority [EFSA]. Scientific opinion on the scrapie situation in the EU after 10 years of monitoring and control in sheep and goats. *EFSA J.* 2014;12(7):3781.
- European Food Safety Authority [EFSA] Scientific Expert Group. Scientific report of the European Food Safety Authority on the evaluation of rapid post mortem TSE tests intended for small ruminants. EFSA; 2005 May. 17 p. Question no. EFSA-Q-2003-084. Available at: http://www.efsa.eu.int/science/tse_assessments/bse_tse/983/biohaz_sr31_smallruminanttsetests_en1.pdf. Accessed 4 Apr. 2007.
- Everest SJ, Thorne L, Barnicle DA, Edwards JC, Elliott H, Jackman R, Hope J. Atypical prion protein in sheep brain collected during the British scrapie-surveillance programme. *J Gen Virol.* 2006;87:471-7.
- Everest SJ, Thorne LT, Hawthorn JA, Jenkins R, Hammersley C, Ramsay AM, Hawkins SA, Venables L, Flynn L, Sayers R, Kilpatrick J, Sach A, Hope J, Jackman R. No abnormal prion protein detected in the milk of cattle infected with the bovine spongiform encephalopathy agent. *J Gen Virol.* 2006;87:2433-41.
- Fast C, Keller M, Balkema-Buschmann A, Hills B, Groschup MH. Complementary studies detecting classical bovine spongiform encephalopathy infectivity in jejunum, ileum and ileocaecal junction in incubating cattle. *Vet Res.* 2013;44:123.
- Fediaevsky A, Gasqui P, Calavas D, Ducrot C. Discrepant epidemiological patterns between classical and atypical scrapie in sheep flocks under French TSE control measures. *Vet J.* 2010;185(3):338-40.
- Fediaevsky A, Tongue SC, Nöremark M, Calavas D, Ru G, Hopp P. A descriptive study of the prevalence of atypical and classical scrapie in sheep in 20 European countries. *BMC Vet Res.* 2008;4:19.
- Fichet G, Comoy E, Dehen C, Challier L, Antloga K, Deslys JP, McDonnell G. Investigations of a prion infectivity assay to evaluate methods of decontamination. *J Microbiol Methods.* 2007;70:511-8.
- Fichet G, Comoy E, Duval C, Antloga K, Dehen C, Charbonnier A, McDonnell G, Brown P, Lasmézas CI, Deslys JP. Novel methods for disinfection of prion-contaminated medical devices. *Lancet.* 2004;364:521-6.
- Fischer JR, Nettles VF (College of Veterinary Medicine, The University of Georgia). National chronic wasting disease surveillance in free-ranging cervids: accomplishments and needs [online]. In: USAHA 2002 Proceedings; 2002 Oct 20-23; St. Louis, MO. Available at: <http://www.usaha.org/speeches/speech02/s02cwwss.html>. * Accessed 2 Dec 2003.
- Foster JD, Goldmann W, Hunter N. Evidence in sheep for pre-natal transmission of scrapie to lambs from infected mothers. *PLoS One.* 2013;8(11):e79433.
- Foster J, McKenzie C, Parnham D, Drummond D, Chong A, Goldman W, Hunter N. Lateral transmission of natural scrapie to scrapie-free New Zealand sheep placed in an endemically infected UK flock. *Vet Rec.* 2006;159:633-4.
- Fragkiadaki EG, Vaccari G, Ekateriniadou LV, Agrimi U, Giadinis ND, Chiappini B, Esposito E, Conte M, Nonno R. PRNP genetic variability and molecular typing of natural goat scrapie isolates in a high number of infected flocks. *Vet Res.* 2011;42:104.
- Franz M, Eiden M, Balkema-Buschmann A, Greenlee J, Schatzl H, Fast C, Richt J, Hildebrandt JP, Groschup MH. Detection of PrP(Sc) in peripheral tissues of clinically affected cattle after oral challenge with bovine spongiform encephalopathy. *J Gen Virol.* 2012;93(Pt 12):2740-8.
- Garza MC, Fernández-Borges N, Bolea R, Badiola JJ, Castilla J, Monleón E. Detection of PrPres in genetically susceptible fetuses from sheep with natural scrapie. *PLoS One.* 2011;6(12):e27525.
- Gavier-Widen D, Noremark M, Benestad S, Simmons M, Renstrom L, Bratberg B, Elvander M, af Segerstad CH. Recognition of the Nor98 variant of scrapie in the Swedish sheep population. *J Vet Diagn Invest.* 2004;16:562-7.
- Gavier-Widén D, Stack MJ, Baron T, Balachandran A, Simmons M. Diagnosis of transmissible spongiform encephalopathies in animals: a review. *J Vet Diagn Invest.* 2005;17:509-27.

Transmissible Spongiform Encephalopathies

- Georgsson G, Sigurdarson S, Brown P. Infectious agent of sheep scrapie may persist in the environment for at least 16 years. *J Gen Virol.* 2006;87:3737-40.
- Gibbs CJ Jr, Amyx HL, Bacote A, Masters CL, Gajdusek DC. Oral transmission of kuru, Creutzfeldt-Jakob disease, and scrapie to nonhuman primates. *J Infect Dis.* 1980;142(2):205-8.
- Gilch S. Overview of chronic wasting disease. In: Kahn CM, Line S, Aiello SE, editors.. *The Merck veterinary manual* [online]. Whitehouse Station, NJ: Merck and Co; 2013. Available at: http://www.merckvetmanual.com/mvm/nervous_system/chronic_wasting_disease/overview_of_chronic_wasting_disease.html. Accessed 20 Jul 2016.
- Gilch S, Chitoor N, Taguchi Y, Stuart M, Jewell JE, Schätzl HM. Chronic wasting disease. *Top Curr Chem.* 2011;305:51-77.
- Giles K, Glidden DV, Beckwith R, Seoanes R, Peretz D, DeArmond SJ, Prusiner SB. Resistance of bovine spongiform encephalopathy (BSE) prions to inactivation. *PLoS Pathog.* 2008 Nov;4(11):e1000206.
- Gill ON, Spencer Y, Richard-Loendt A, Kelly C, Dabaghian R, et al. Prevalent abnormal prion protein in human appendixes after bovine spongiform encephalopathy epizootic: large scale survey. *BMJ.* 2013 Oct 15;347:f5675.
- Giovannini A, Savini L, Conte A, Fiore GL. Comparison of BSE prevalence estimates from EU countries for the period July to December 2001 to the OIE and EU GBR classifications. *J Vet Med B Infect Dis Vet Public Health.* 2005;52:262-71.
- Goldmann W. PrP genetics in ruminant transmissible spongiform encephalopathies. *Vet Res.* 2008;39:30.
- Goldmann W, Hunter N, Smith G, Foster J, Hope J. PrP genotype and agent effects in scrapie: change in allelic interaction with different isolates of agent in sheep, a natural host of scrapie. *J Gen Virol.* 1994;75:989-95.
- González L, Chianini F, Martin S, Sisó S, Gibbard L, Reid HW, Jeffrey M. Comparative titration of experimental ovine BSE infectivity in sheep and mice. *J Gen Virol.* 2007;88:714-7.
- González L, Dagleish MP, Martin S, Dexter G, Steele P, Finlayson J, Jeffrey M. Diagnosis of preclinical scrapie in live sheep by the immunohistochemical examination of rectal biopsies. *Vet Rec.* 2008;162(13):397-403.
- González L, Martin S, Sisó S, Konold T, Ortiz-Peláez A, Phelan L, Goldmann W, Stewart P, Saunders G, Windl O, Jeffrey M, Hawkins SA, Dawson M, Hope J. High prevalence of scrapie in a dairy goat herd: tissue distribution of disease-associated PrP and effect of PRNP genotype and age. *Vet Res.* 2009;40(6):65.
- González L, Pitarch JL, Martin S, Thurston L, Moore J, Acín C, Jeffrey M. Identical pathogenesis and neuropathological phenotype of scrapie in valine, arginine, glutamine/valine, arginine, glutamine sheep infected experimentally by the oral and conjunctival routes. *J Comp Pathol.* 2014;150(1):47-56.
- Gonzalez L, Pitarch JL, Martin S, Thurston L, Simmons H, Acin C, Jeffrey M. Influence of polymorphisms in the prion protein gene on the pathogenesis and neuropathological phenotype of sheep scrapie after oral infection. *J Comp Pathol.* 2014;150:57-70.
- Götte DR, Benestad SL, Laude H, Zurbriggen A, Oevermann A, Seuberlich T. Atypical scrapie isolates involve a uniform prion species with a complex molecular signature. *PLoS One.* 2011;6(11):e27510.
- Gough KC, Baker CA, Rees HC, Terry LA, Spiropoulos J, Thorne L, Maddison BC. The oral secretion of infectious scrapie prions occurs in preclinical sheep with a range of PRNP genotypes. *J Virol.* 2012;86(1):566-71.
- Gough KC, Baker CA, Simmons HA, Hawkins SA, Maddison BC. Circulation of prions within dust on a scrapie affected farm. *Vet Res.* 2015;46:40.
- Gough KC, Bishop K, Maddison BC. Highly sensitive detection of small ruminant bovine spongiform encephalopathy within transmissible spongiform encephalopathy mixes by serial protein misfolding cyclic amplification. *J Clin Microbiol.* 2014;52(11):3863-8.
- Gough KC, Rees HC, Ives SE, Maddison BC. Methods for differentiating prion types in food-producing animals. *Biology (Basel).* 2015;4(4):785-813.
- Gough KC, Maddison BC. Prion transmission: prion excretion and occurrence in the environment. *Prion.* 2010;4(4):275-82.
- Gould DH, Voss JL, Miller MW, Bachand AM, Cummings BA, Frank AA. Survey of cattle in northeast Colorado for evidence of chronic wasting disease: geographical and high-risk targeted sample. *J Vet Diagn Invest.* 2003;15:274-7.
- Gray JG, Dudas S, Graham C, Czub S. Performance analysis of rapid diagnostic tests on atypical bovine spongiform encephalopathy. *J Vet Diagn Invest.* 2012;24(5):976-80.
- Greenwood P. Federal disease control--scrapie. *Can Vet J.* 2002;43:625-9.
- Hadlow WJ, Karstad L. Transmissible encephalopathy of mink in Ontario. *Can Vet J.* 1968;9:193-6.
- Haley NJ, Hoover EA. Chronic wasting disease of cervids: current knowledge and future perspectives. *Annu Rev Anim Biosci.* 2015;3:305-25.
- Haley NJ, Mathiason CK, Carver S, Zabel M, Telling GC, Hoover EA. Detection of chronic wasting disease prions in salivary, urinary, and intestinal tissues of deer: potential mechanisms of prion shedding and transmission. *J Virol.* 2011;85(13):6309-18.
- Haley NJ, Mathiason CK, Zabel MD, Telling GC, Hoover EA. Detection of sub-clinical CWD infection in conventional test-negative deer long after oral exposure to urine and feces from CWD+ deer. *PLoS One.* 2009;4(11):e7990.
- Haley NJ, Seelig DM, Zabel MD, Telling GC, Hoover EA. Detection of CWD prions in urine and saliva of deer by transgenic mouse bioassay. *PLoS One.* 2009;4(3):e4848.
- Hall V, Brookes D, Nacul L, Gill ON, Connor N; CJD Incidents Panel. Managing the risk of iatrogenic transmission of Creutzfeldt-Jakob disease in the UK. *J Hosp Infect.* 2014;88(1):22-7.
- Hamir AN, Clark WW, Sutton DL, Miller JM, Stack MJ, Chaplin MJ, Jenny AL. Resistance of domestic cats to a US sheep scrapie agent by intracerebral route. *J Vet Diagn Invest.* 2002;14(5):444-5.
- Hamir AN, Miller JM, O'Rourke KI, Bartz JC, Stack MJ, Chaplin MJ. Transmission of transmissible mink encephalopathy to raccoons (*Procyon lotor*) by intracerebral inoculation. *J Vet Diagn Invest.* 2004;16:57-63.
- Harrington RD, Baszler TV, O'Rourke KI, Schneider DA, Spraker TR, Liggitt HD, Knowles DP. A species barrier limits transmission of chronic wasting disease to mink (*Mustela vison*). *J Gen Virol.* 2008;89(Pt 4):1086-96.

Transmissible Spongiform Encephalopathies

- Hawkins SA, Simmons HA, Gough KC, Maddison BC. Persistence of ovine scrapie infectivity in a farm environment following cleaning and decontamination. *Vet Rec*. 2015;176(4):99.
- Henderson DM, Davenport KA, Haley NJ, Denkers ND, Mathiason CK, Hoover EA. Quantitative assessment of prion infectivity in tissues and body fluids by real-time quaking-induced conversion. *J Gen Virol*. 2015;96(Pt 1):210-9.
- Henderson DM, Denkers ND, Hoover CE, Garbino N, Mathiason CK, Hoover EA. Longitudinal detection of prion shedding in saliva and urine by chronic wasting disease-infected deer by real-time quaking-induced conversion. *J Virol*. 2015;89(18):9338-47.
- Henderson DM, Manca M, Haley NJ, Denkers ND, Nalls AV, Mathiason CK, Caughey B, Hoover EA. Rapid antemortem detection of CWD prions in deer saliva. *PLoS One*. 2013;8(9):e74377.
- Henry C, Knight R. Clinical features of variant Creutzfeldt-Jakob disease. *Rev Med Virol*. 2002;12:143-50.
- Hilbe MM, Soldati GG, Zlinszky KK, Wunderlin SS, Ehrensperger FF. Immunohistochemical study of PrP(Sc) distribution in neural and extraneural tissues of two cats with feline spongiform encephalopathy. *BMC Vet Res*. 2009;5:11.
- Hildebrandt H. Transmissible mink encephalopathy In: Kahn CM, Line S, Aiello SE, editors. *The Merck veterinary manual* [online]. Whitehouse Station, NJ: Merck and Co; 2014. . Available at: http://www.merckvetmanual.com/mvm/exotic_and_laboratory_animals/mink/prion_disease_of_mink.html. Accessed 1 Sept 2016.
- Hill AF, Collinge J. Subclinical prion infection in humans and animals. *Br Med Bull*. 2003;66:161-70.
- Hilton DA. Pathogenesis and prevalence of variant Creutzfeldt-Jakob disease. *J Pathol*. 2006;208:134-41.
- Hirata Y, Ito H, Furuta T, Ikuta K, Sakudo A. Degradation and destabilization of abnormal prion protein using alkaline detergents and proteases. *Int J Mol Med*. 2010;25(2):267-70.
- Hoffmann C, Eiden M, Kaatz M, Keller M, Ziegler U, Rogers R, Hills B, Balkema-Buschmann A, van Keulen L, Jacobs JG, Groschup MH. BSE infectivity in jejunum, ileum and ileocaecal junction of incubating cattle. *Vet Res*. 2011;42(1):21.
- Hoffmann C, Ziegler U, Buschmann A, Weber A, Kupfer L, Oelschlegel A, Hammerschmidt B, Groschup MH. Prions spread via the autonomic nervous system from the gut to the central nervous system in cattle incubating bovine spongiform encephalopathy. *J Gen Virol*. 2007;88:1048-55.
- Hoinville LJ, Tongue SC, Wilesmith JW. Evidence for maternal transmission of scrapie in naturally affected flocks. *Prev Vet Med*. 2010;93(2-3):121-8.
- Horby P. Variant Creutzfeldt-Jakob disease: an unfolding epidemic of misfolded proteins. *J Paediatr Child Health*. 2002;38:539-42.
- Houston F, McCutcheon S, Goldmann W, Chong A, Foster J, Sisó S, González L, Jeffrey M, Hunter N. Prion diseases are efficiently transmitted by blood transfusion in sheep. *Blood*. 2008;112(12):4739-45.
- Hunter N. Scrapie and experimental BSE in sheep. *Br Med Bull*. 2003;66:171-83.
- Irani DN. Johns Hopkins Department of Neurology. Resource on prion diseases [online]. Transmissible mink encephalopathy. Available at: <http://www.jhu-prion.org/animal/anitme-hist.shtml>. * Accessed 7 Nov 2001.
- Iulini B, Cantile C, Mandara MT, Maurella C, Loria GR, Castagnaro M, Salvadori C, Porcario C, Corona C, Perazzini AZ, Maroni A, Caramelli M, Casalone C. Neuropathology of Italian cats in feline spongiform encephalopathy surveillance. *Vet Pathol*. 2008;45(5):626-33.
- Iwamaru Y, Imamura M, Matsuura Y, Masujin K, Shimizu Y, Shu Y, Kurachi M, Kasai K, Murayama Y, Fukuda S, Onoe S, Hagiwara K, Yamakawa Y, Sata T, Mohri S, Okada H, Yokoyama T. Accumulation of L-type bovine prions in peripheral nerve tissues. *Emerg Infect Dis*. 2010;16(7):1151-4.
- Jacques CN, Jenks JA, Jenny AL, Griffin SL. Prevalence of chronic wasting disease and bovine tuberculosis in free-ranging deer and elk in South Dakota. *J Wildl Dis*. 2003;39:29-34.
- Jewell JE, Brown J, Kreeger T, Williams ES. Prion protein in cardiac muscle of elk (*Cervus elaphus nelsoni*) and white-tailed deer (*Odocoileus virginianus*) infected with chronic wasting disease. *J Gen Virol*. 2006;87:3443-50.
- Jennelle CS, Samuel MD, Nolden CA, Keane DP, Barr DJ, Johnson C, Vanderloo JP, Aiken JM, Hamir AN, Hoover EA. Surveillance for transmissible spongiform encephalopathy in scavengers of white-tailed deer carcasses in the chronic wasting disease area of Wisconsin. *J Toxicol Environ Health A*. 2009;72(17-18):1018-24.
- Kanata E, Humphreys-Panagiotidis C, Giadinis ND, Papaioannou N, Arsenakis M, Sklavadias T. Perspectives of a scrapie resistance breeding scheme targeting Q211, S146 and K222 caprine PRNP alleles in Greek goats. *Vet Res*. 2014;45:43.
- Kariv-Inbal Z, Ben-Hur T, Grigoriadis NC, Engelstein R, Gabizon R. Urine from scrapie-infected hamsters comprises low levels of prion infectivity. *Neurodegener Dis*. 2006;3:123-8.
- Keane D, Barr D, Osborn R, Langenberg J, O'Rourke K, Schneider D, Bochsler P. Validation of use of rectoanal mucosa-associated lymphoid tissue for immunohistochemical diagnosis of chronic wasting disease in white-tailed deer (*Odocoileus virginianus*). *J Clin Microbiol*. 2009;47(5):1412-7.
- Kelly DF, Wells GA, Haritani M, Higgins RJ, Jeffrey M. Neuropathological findings in cats with clinically suspect but histologically unconfirmed feline spongiform encephalopathy. *Vet Rec*. 2005;156:472-7.
- Kim TY, Shon HJ, Joo YS, Mun UK, Kang KS, Lee YS. Additional cases of chronic wasting disease in imported deer in Korea. *J Vet Med Sci*. 2005;67:753-9.
- Kimura K, Haritani M. Distribution of accumulated prion protein in a cow with bovine spongiform encephalopathy. *Vet Rec*. 2008;162(25):822-5.
- Kittelberger R, Chaplin MJ, Simmons MM, Ramirez-Villaescusa A, McIntyre L, MacDiarmid SC, Hannah MJ, Jenner J, Bueno R, Bayliss D, Black H, Pigott CJ, O'Keefe JS. Atypical scrapie/Nor98 in a sheep from New Zealand. *J Vet Diagn Invest*. 2010;22(6):863-75.

Transmissible Spongiform Encephalopathies

- Kong Q, Huang S, Zou W, Vanegas D, Wang M, Wu D, Yuan J, Zheng M, Bai H, Deng H, Chen K, Jenny AL, O'Rourke K, Belay ED, Schonberger LB, Petersen RB, Sy MS, Chen SG, Gambetti P. Chronic wasting disease of elk: transmissibility to humans examined by transgenic mouse models. *J Neurosci*. 2005;25:7944-9.
- Konold T, Bone GE, Phelan LJ, Simmons MM, González L, Sisó S, Goldmann W, Cawthraw S, Hawkins SA. Monitoring of clinical signs in goats with transmissible spongiform encephalopathies. *BMC Vet Res*. 2010 4;6:13.
- Konold T, Bone G, Ryder S, Hawkins SA, Courtin F, Berthelin-Baker C. Clinical findings in 78 suspected cases of bovine spongiform encephalopathy in Great Britain. *Vet Rec*. 2004;155:659-66.
- Konold T, Davis A, Bone G, Bracegirdle J, Everitt S, Chaplin M, Saunders GC, Cawthraw S, Simmons MM. Clinical findings in two cases of atypical scrapie in sheep: a case report. *BMC Vet Res*. 2007;13:3:2.
- Konold T, Moore SJ, Bellworthy SJ, Simmons HA. Evidence of scrapie transmission via milk. *BMC Vet Res*. 2008;4:14.
- Konold T, Moore SJ, Bellworthy SJ, Terry LA, Thorne L, Ramsay A, Salguero FJ, Simmons MM, Simmons HA. Evidence of effective scrapie transmission via colostrum and milk in sheep. *BMC Vet Res*. 2013;9:99.
- Konold T, Phelan L. Clinical examination protocol for the detection of scrapie. *Vet Rec*. 2014;174(10):257.
- Konold T, Simmons HA, Webb PR, Bellerby PJ, Hawkins SA, González L. Transmission of classical scrapie via goat milk. *Vet Rec*. 2013;172(17):455.
- Kreeger TJ, Montgomery DL, Jewell JE, Schultz W, Williams ES. Oral transmission of chronic wasting disease in captive Shira's moose. *J Wildl Dis*. 2006;42:640-5.
- Kubler E, Oesch B, Raeber AJ. Diagnosis of prion diseases. *Br Med Bull*. 2003;66:267-79.
- Kurt TD, Sigurdson CJ. Cross-species transmission of CWD prions. *Prion*. 2016;10(1):83-91.
- Lacroux C, Comoy E, Moudjou M, Perret-Liaudet A, Lugan S, et al. Preclinical detection of variant CJD and BSE prions in blood. *PLoS Pathog*. 2014;10(6):e1004202.
- Lacroux C, Corbiere F, Tabouret G, Lugan S, Costes P, Mathey J, Delmas JM, Weisbecker JL, Foucras G, Cassard H, Elsen JM, Schelcher F, Andreoletti O. Dynamics and genetics of PrPSc placental accumulation in sheep. *J Gen Virol*. 2007;88:1056-61.
- Lacroux C, Perrin-Chauvineau C, Corbière F, Aron N, Aguilar-Calvo P, Torres JM, Costes P, Brémaud I, Lugan S, Schelcher F, Barillet F, Andreoletti O. Genetic resistance to scrapie infection in experimentally challenged goats. *J Virol*. 2014;88(5):2406-13.
- Lacroux C, Simon S, Benestad SL, Mailliet S, Mathey J, et al. Prions in milk from ewes incubating natural scrapie. *PLoS Pathog*. 2008;4:1000238.
- Lasmézas CI, Comoy E, Hawkins S, Herzog C, Mouthon F, Konold T, Auvré F, Correia E, Lescoutra-Etcheagaray N, Salès N, Wells G, Brown P, Deslys JP. Risk of oral infection with bovine spongiform encephalopathy agent in primates. *Lancet*. 2005;365:781-3.
- Le Dur A, Beringue V, Andreoletti O, Reine F, Lai TL, Baron T, Bratberg B, Vilotte JL, Sarradin P, Benestad SL, Laude H. A newly identified type of scrapie agent can naturally infect sheep with resistant PrP genotypes. *Proc Natl Acad Sci U S A*. 2005;102:16031-6.
- Lehmann S, Pastore M, Rogez-Kreuz C, Richard M, Belondrade M, Rauwel G, Durand F, Yousfi R, Criquelion J, Clayette P, Perret-Liaudet A. New hospital disinfection processes for both conventional and prion infectious agents compatible with thermosensitive medical equipment. *J Hosp Infect*. 2009;72(4):342-50.
- Lezmi S, Bencsik A, Monks E, Petit T, Baron T. First case of feline spongiform encephalopathy in a captive cheetah born in France: PrP(sc) analysis in various tissues revealed unexpected targeting of kidney and adrenal gland. *Histochem Cell Biol*. 2003;119:415-22.
- Lezmi S, Baron TG, Bencsik AA. Is the presence of abnormal prion protein in the renal glomeruli of feline species presenting with FSE authentic? *BMC Vet Res*. 2010;6:41.
- Liberski PP, Sikorska B, Guiryo D, Bessen RA. Transmissible mink encephalopathy - review of the etiology of a rare prion disease. *Folia Neuropathol*. 2009;47(2):195-204.
- Lord Phillips, chair. The BSE inquiry: The report. A report to the Minister of Agriculture, Fisheries and Food, the Secretary of State for Health and the Secretaries of State for Scotland, Wales and Northern Ireland. Report no. HC 887-1. London: Her Majesty's Stationery Office; 2000. Available at: <http://www.bseinquiry.gov.uk/report/>. Accessed 2006 Jan.
- Ludlam CA, Turner ML. Managing the risk of transmission of variant Creutzfeldt Jakob disease by blood products. *Br J Haematol*. 2006;132:13-24.
- Luhken G, Buschmann A, Brandt H, Eiden M, Groschup MH, Erhardt G. Epidemiological and genetical differences between classical and atypical scrapie cases. *Vet Res*. 2007;38:65-80.
- Maddison BC, Baker CA, Rees HC, Terry LA, Thorne L, Bellworthy SJ, Whitlam GC, Gough KC. Prions are secreted in milk from clinically normal scrapie-exposed sheep. *J Virol*. 2009;83(16):8293-6.
- Maddison BC, Baker CA, Terry LA, Bellworthy SJ, Thorne L, Rees HC, Gough KC. Environmental sources of scrapie prions. *J Virol*. 2010;84(21):11560-2.
- Maddison BC, Owen JP, Bishop K, Shaw G, Rees HC, Gough KC. The interaction of ruminant PrP(Sc) with soils is influenced by prion source and soil type. *Environ Sci Technol*. 2010;44(22):8503-8.
- Maddison BC, Rees HC, Baker CA, Taema M, Bellworthy SJ, Thorne L, Terry LA, Gough KC. Prions are secreted into the oral cavity in sheep with preclinical scrapie. *J Infect Dis*. 2010;201(11):1672-6.
- Maestrle C, Cancedda MG, Pintus D, Masia M, Nonno R, Ru G, Carta A, Demontis F, Santucci C, Ligios C. Genetic and pathological follow-up study of goats experimentally and naturally exposed to a sheep scrapie isolate. *J Virol*. 2015;89(19):10044-52.
- Manjerovic MB, Green ML, Mateus-Pinilla N, Novakofski J. The importance of localized culling in stabilizing chronic wasting disease prevalence in white-tailed deer populations. *Prev Vet Med*. 2014;113(1):139-45.

Transmissible Spongiform Encephalopathies

- Marsh RF. Animal model of human disease: Kuru, Creutzfeldt-Jakob disease (slow virus infections). Animal model: transmissible mink encephalopathy, scrapie-like disease of mink. *Am J Pathol.* 1972;69:209-12.
- Marsh RF, Kincaid AE, Bessen RA, Bartz JC. Interspecies transmission of chronic wasting disease prions to squirrel monkeys (*Saimiri sciureus*). *J Virol.* 2005;79:13794-6.
- Martin S, Jeffrey M, González L, Sisó S, Reid HW, Steele P, Dagleish MP, Stack MJ, Chaplin MJ, Balachandran A. Immunohistochemical and biochemical characteristics of BSE and CWD in experimentally infected European red deer (*Cervus elaphus elaphus*). *BMC Vet Res.* 2009;5:26.
- Masujin K, Matthews D, Wells GA, Mohri S, Yokoyama T. Prions in the peripheral nerves of bovine spongiform encephalopathy-affected cattle. *J Gen Virol.* 2007;88:1850-8.
- Masujin K, Orrú CD, Miyazawa K, Groveman BR, Raymond LD, Hughson AG, Caughey B. Detection of atypical H-type bovine spongiform encephalopathy and discrimination of bovine prion strains by real-time quaking-induced conversion. *J Clin Microbiol.* 2016;54(3):676-86.
- Mathiason CK, Hays SA, Powers J, Hayes-Klug J, Langenberg J, Dahmes SJ, Osborn DA, Miller KV, Warren RJ, Mason GL, Hoover EA. Infectious prions in pre-clinical deer and transmission of chronic wasting disease solely by environmental exposure. *PLoS One.* 2009;4(6):e5916.
- Mathiason CK, Powers JG, Dahmes SJ, Osborn DA, Miller KV, et al. Infectious prions in the saliva and blood of deer with chronic wasting disease. *Science.* 2006;314:133-6.
- Matsuura Y, Ishikawa Y, Bo X, Murayama Y, Yokoyama T, Somerville RA, Kitamoto T, Mohri S. Quantitative analysis of wet-heat inactivation in bovine spongiform encephalopathy. *Biochem Biophys Res Commun.* 2013;432(1):86-91.
- McDonnell G, Dehen C, Perrin A, Thomas V, Igel-Egalon A, Burke PA, Deslys JP, Comoy E. Cleaning, disinfection and sterilization of surface prion contamination. *J Hosp Infect.* 2013;85(4):268-73.
- McKenzie D, Bartz JC, Marsh RF. Transmissible mink encephalopathy. *Semin Virol.* 1996;7:201-6.
- Meloni D, Davidse A, Langeveld JP, Varello K, Casalone C, Corona C, Balkema-Buschmann A, Groschup MH, Ingravalle F, Bozzetta E. EU-approved rapid tests for bovine spongiform encephalopathy detect atypical forms: a study for their sensitivities. *PLoS One.* 2012;7(9):e43133.
- Miller MW, Wild MA. Epidemiology of chronic wasting disease in captive white-tailed and mule deer. *J Wildl Dis.* 2004;40:320-7.
- Miller MW, Williams ES, Hobbs NT, Wolfe LL. Environmental sources of prion transmission in mule deer. *Emerg Infect Dis.* 2004;10:1003-6.
- Monello RJ, Powers JG, Hobbs NT, Spraker TR, O'Rourke KI, Wild MA. Efficacy of antemortem rectal biopsies to diagnose and estimate prevalence of chronic wasting disease in free-ranging cow elk (*Cervus elaphus nelsoni*). *Wildl Dis.* 2013 Apr;49(2):270-8.
- Monleón E, Garza MC, Sarasa R, Alvarez-Rodríguez J, Bolea R, Monzón M, Vargas MA, Badiola JJ, Acín C. An assessment of the efficiency of PrPsc detection in rectal mucosa and third-eyelid biopsies from animals infected with scrapie. *Vet Microbiol.* 2011;147(3-4):237-43.
- Morawski AR, Carlson CM, Chang H, Johnson CJ. *In vitro* prion protein conversion suggests risk of bighorn sheep (*Ovis canadensis*) to transmissible spongiform encephalopathies. *BMC Vet Res.* 2013;9:157.
- Moum T, Olsaker I, Hopp P, Moldal T, Valheim M, Moum T, Benestad SL. Polymorphisms at codons 141 and 154 in the ovine prion protein gene are associated with scrapie Nor98 cases. *J Gen Virol.* 2005;86:231-5.
- Nalls AV, McNulty E, Powers J, Seelig DM, Hoover C, Haley NJ, Hayes-Klug J, Anderson K, Stewart P, Goldmann W, Hoover EA, Mathiason CK. Mother to offspring transmission of chronic wasting disease in reeves' muntjac deer. *PLoS One.* 2013;8(8):e71844.
- Nichols TA, Fischer JW, Spraker TR, Kong Q, VerCauteren KC. CWD prions remain infectious after passage through the digestive system of coyotes (*Canis latrans*). *Prion.* 2015;9(5):367-75.
- Nichols TA, Spraker TR, Rigg TD, Meyerrett-Reid C, Hoover C, Michel B, Bian J, Hoover E, Gidlewski T, Balachandran A, O'Rourke K, Telling GC, Bowen R, Zabel MD, VerCauteren KC. Intranasal inoculation of white-tailed deer (*Odocoileus virginianus*) with lyophilized chronic wasting disease prion particulate complexed to montmorillonite clay. *PLoS One.* 2013;8(5):e62455.
- Norwegian Food Safety Authority. Chronic wasting disease in Norway. 2016. Available at: http://www.mattilsynet.no/language/english/animals/chronic_wasting_disease_in_norway.23274. Accessed 28 Jul 2016.
- Novakofski J, Brewer MS, Mateus-Pinilla N, Killefer J, McCusker RH. Prion biology relevant to bovine spongiform encephalopathy. *J Anim Sci.* 2005;83:1455-76.
- Okada H, Iwamaru Y, Fukuda S, Yokoyama T, Mohri S. Detection of disease-associated prion protein in the optic nerve and the adrenal gland of cattle with bovine spongiform encephalopathy by using highly sensitive immunolabeling procedures. *J Histochem Cytochem.* 2012;60(4):290-300.
- Okada H, Miyazawa K, Fukuda S, Iwamaru Y, Imamura M, Masujin K, Matsuura Y, Fujii T, Fujii K, Kageyama S, Yoshioka M, Murayama Y, Yokoyama T. The presence of disease-associated prion protein in skeletal muscle of cattle infected with classical bovine spongiform encephalopathy. *J Vet Med Sci.* 2014;76(1):103-7.
- Okada H, Miyazawa K, Imamura M, Iwamaru Y, Masujin K, Matsuura Y, Yokoyama T. Transmission of atypical scrapie to homozygous ARQ sheep. *J Vet Med Sci.* 2016 Jun 20. [Epub ahead of print]
- Olszowy KM, Lavelle J, Rachfal K, Hempstead S, Drouin K, Darcy JM 2nd, Reiber C, Garruto RM. Six-year follow-up of a point-source exposure to CWD contaminated venison in an Upstate New York community: risk behaviours and health outcomes 2005-2011. *Public Health.* 2014;128(9):860-8.
- Onnasch H, Gunn HM, Bradshaw BJ, Benestad SL, Bassett HF. Two Irish cases of scrapie resembling Nor98. *Vet Rec.* 2004;155:636-7.
- Orge L, Galo A, Machado C, Lima C, Ochoa C, Silva J, Ramos M, Simas JP. Identification of putative atypical scrapie in sheep in Portugal. *J Gen Virol.* 2004;85:3487-3491.
- Orge L, Machado CG, Ramalho L, Carvalho R, Silva J, Almeida P, Tavares P, Ochoa C, Lima C, Pinto MJ, Simas JP. Identification of H-type BSE in Portugal. *Prion.* 2015;9(1):22-8.

Transmissible Spongiform Encephalopathies

- Orrú CD, Favole A, Corona C, Mazza M, Manca M, Groveman BR, Hughson AG, Acutis PL, Caramelli M, Zanusso G, Casalone C, Caughey B. Detection and discrimination of classical and atypical L-type bovine spongiform encephalopathy by real-time quaking-induced conversion. *Clin Microbiol*. 2015;53(4):1115-20.
- Pattison J. The emergence of bovine spongiform encephalopathy and related diseases. *Emerg Infect Dis*. 1998;4:390-4.
- Pirisinu L, Migliore S, Di Bari MA, Esposito E, Baron T, D'Agostino C, De Grossi L, Vaccari G, Agrimi U, Nonno R. Molecular discrimination of sheep bovine spongiform encephalopathy from scrapie. *Emerg Infect Dis*. 2011;17(4):695-8.
- Promed Mail. Chronic wasting disease, elk - South Korea (Kyungsang). Nov 25, 2004. Archive Number 20041125.3155. Available at **Error! Hyperlink reference not valid.** Accessed 30 Sept 2008.
- Promed Mail. Chronic wasting disease, cervid - Europe (02): (Norway) moose. Archive Number: 20160614.4276448 Available at: **Error! Hyperlink reference not valid.** Accessed 28 Jul 2016.
- Promed Mail. Feline spongiform encephalopathy, cat – Switzerland. Aug 24, 2003. Archive Number 20030824.2132. Available at **Error! Hyperlink reference not valid.** Accessed 19 Sept 2007.
- Promed Mail. Feline spongiform encephalopathy, cheetah – Germany. Sept 13, 2007. Archive Number 20070913.3038. Available at <http://www.promedmail.org>. Accessed 19 Sept 2007.
- Pulford B, Spraker TR, Wyckoff AC, Meyerett C, Bender H, Ferguson A, Wyatt B, Lockwood K, Powers J, Telling GC, Wild MA, Zabel MD. Detection of PrPCWD in feces from naturally exposed Rocky Mountain elk (*Cervus elaphus nelsoni*) using protein misfolding cyclic amplification. *J Wildl Dis*. 2012;48(2):425-34.
- Race B, Meade-White KD, Miller MW, Barbian KD, Rubenstein R, et al. Susceptibilities of nonhuman primates to chronic wasting disease. *Emerg Infect Dis*. 2009;15(9):1366-76.
- Race B, Meade-White KD, Phillips K, Striebel J, Race R, Chesebro B. Chronic wasting disease agents in nonhuman primates. *Emerg Infect Dis*. 2014;20(5):833-7.
- Race BL, Meade-White KD, Ward A, Jewell J, Miller MW, Williams ES, Chesebro B, Race RE. Levels of abnormal prion protein in deer and elk with chronic wasting disease. *Emerg Infect Dis*. 2007;13:824-30.
- Raymond GJ, Raymond LD, Meade-White KD, Hughson AG, Favara C, Gardner D, Williams ES, Miller MW, Race RE, Caughey B. Transmission and adaptation of chronic wasting disease to hamsters and transgenic mice: evidence for strains. *J Virol*. 2007;81:4305-14.
- Richt JA, Kunkle RA, Alt D, Nicholson EM, Hamir AN, Czub S, Kluge J, Davis AJ, Hall SM. Identification and characterization of two bovine spongiform encephalopathy cases diagnosed in the United States. *J Vet Diagn Invest*. 2007;19:142-54.
- Robinson MM, Hadlow WJ, Huff TP, Wells GA, Dawson M, Marsh RF, Gorham JR. Experimental infection of mink with bovine spongiform encephalopathy. *J Gen Virol*. 1994;75:2151-5.
- Robinson SJ, Samuel MD, O'Rourke KI, Johnson CJ. The role of genetics in chronic wasting disease of North American cervids. *Prion*. 2012;6(2):153-62.
- Rodríguez-Martínez AB, Garrido JM, Maza S, Benedicto L, Geijo M, Gómez N, Minguijón E, Benestad SL, Juste RA. Atypical/Nor98 scrapie in the Basque country: a case report of eight outbreaks. *BMC Vet Res*. 2010;6:17.
- Rogez-Kreuz C, Yousfi R, Soufflet C, Quadrio I, Yan ZX, Huyot V, Aubenque C, Destrez P, Roth K, Roberts C, Favero M, Clayette P. Inactivation of animal and human prions by hydrogen peroxide gas plasma sterilization. *Infect Control Hosp Epidemiol*. 2009;30(8):769-77.
- Ronzon F, Bencsik A, Lezmi S, Vulin J, Kodjo A, Baron T. BSE inoculation to prion diseases-resistant sheep reveals tricky silent carriers. *Biochem Biophys Res Commun*. 2006;350:872-7.
- Rubenstein R, Chang B, Gray P, Piltch M, Bulgin MS, Sorensen-Melson S, Miller MW. Prion disease detection, PMCA kinetics, and IgG in urine from sheep naturally/experimentally infected with scrapie and deer with preclinical/clinical chronic wasting disease. *J Virol*. 2011;85(17):9031-8.
- Rutala WA, Weber DJ; Society for Healthcare Epidemiology of America. Guideline for disinfection and sterilization of prion-contaminated medical instruments. *Infect Control Hosp Epidemiol*. 2010;31(2):107-17.
- Ryder S, Dexter G, Bellworthy S, Tongue S. Demonstration of lateral transmission of scrapie between sheep kept under natural conditions using lymphoid tissue biopsy. *Res Vet Sci*. 2004;76:211-7.
- Saba R, Booth SA. The genetics of susceptibility to variant Creutzfeldt-Jakob disease. *Public Health Genomics*. 2013;16(1-2):17-24.
- Salta E, Panagiotidis C, Teliousis K, Petrakis S, Eleftheriadis E, Arapoglou F, Grigoriadis N, Nicolaou A, Kaldrymidou E, Krey G, Sklaviadis T. Evaluation of the possible transmission of BSE and scrapie to gilthead sea bream (*Sparus aurata*). *PLoS One*. 2009 Jul 28;4(7):e6175.
- Saunders SE, Bartelt-Hunt SL, Bartz JC. Prions in the environment: occurrence, fate and mitigation. *Prion*. 2008;2(4):162-9.
- Saunders SE, Bartz JC, Telling GC, Bartelt-Hunt SL. Environmentally-relevant forms of the prion protein. *Environ Sci Technol*. 2008;42(17):6573-9.
- Saunders SE, Bartz JC, Vercauteren KC, Bartelt-Hunt SL. An enzymatic treatment of soil-bound prions effectively inhibits replication. *Appl Environ Microbiol*. 2011;77(13):4313-7.
- Schneider DA, Harrington RD, Zhuang D, Yan H, Truscott TC, Dassanayake RP, O'Rourke KI. Disease-associated prion protein in neural and lymphoid tissues of mink (*Mustela vison*) inoculated with transmissible mink encephalopathy. *J Comp Pathol*. 2012;147(4):508-21.
- Schneider DA, Madsen-Bouterse SA, Zhuang D, Truscott TC, Dassanayake RP, O'Rourke KI. The placenta shed from goats with classical scrapie is infectious to goat kids and lambs. *J Gen Virol*. 2015;96(8):2464-9.
- Schuler KL, Jenks JA, DePerno CS, Wild MA, Swanson CC. Tonsillar biopsy test for chronic wasting disease: Two sampling approaches in mule deer and white-tailed deer. *J Wildl Dis*. 2005;41:820-4.
- Schwabenlander MD, Culhane MR, Hall SM, Goyal SM, Anderson PL, Carstensen M, Wells SJ, Slade WB, Armién AG. A case of chronic wasting disease in a captive red deer (*Cervus elaphus*). *J Vet Diagn Invest*. 2013;25(5):573-6.

Transmissible Spongiform Encephalopathies

- Seeger H, Heikenwalder M, Zeller N, Kranich J, Schwarz P, Gaspert A, Seifert B, Miele G, Aguzzi A. Coincident scrapie infection and nephritis lead to urinary prion excretion. *Science*. 2005;310:324-6.
- Seidel B, Thomzig A, Buschmann A, Groschup MH, Peters R, Beekes M, Terytze K. Scrapie agent (Strain 263K) can transmit disease via the oral route after persistence in soil over years. *PLoS ONE*. 2007;2(5):e435.
- Selariu A, Powers JG, Nalls A, Brandhuber M, Mayfield A, Fullaway S, Wyckoff CA, Goldmann W, Zabel MM, Wild MA, Hoover EA, Mathiason CK. *In utero* transmission and tissue distribution of chronic wasting disease-associated prions in free-ranging Rocky Mountain elk. *J Gen Virol*. 2015;96(11):3444-55.
- Seuberlich T. Overview of bovine encephalopathy. Kahn CM, Line S, Aiello SE, editors. *The Merck veterinary manual* [online]. Whitehouse Station, NJ: Merck and Co; 2014. Bovine spongiform encephalopathy. Available at: http://www.merckvetmanual.com/mvm/nervous_system/bovine_spongiform_encephalopathy/overview_of_bovine_spongiform_encephalopathy.html. Accessed 26 Aug 2016.
- Seuberlich T, Heim D, Zurbriggen A. Atypical transmissible spongiform encephalopathies in ruminants: a challenge for disease surveillance and control. *J Vet Diagn Invest*. 2010;22(6):823-42.
- Shimada K, Hayashi HK, Ookubo Y, Iwamaru Y, Imamura M, Takata M, Schmerr MJ, Shinagawa M, Yokoyama T. Rapid PrP(Sc) detection in lymphoid tissue and application to scrapie surveillance of fallen stock in Japan: variable PrP(Sc) accumulation in palatal tonsil in natural scrapie. *Microbiol Immunol*. 2005;49:801-4.
- Sigurdson CJ, Aguzzi A. Chronic wasting disease. *Biochim Biophys Acta*. 2007;1772:610-8.
- Sigurdson CJ, Miller MW. Other animal prion diseases. *Br Med Bull*. 2003;66:199-212.
- Simmons MM, Konold T, Simmons HA, Spencer YI, Lockey R, Spiropoulos J, Everitt S, Clifford D. Experimental transmission of atypical scrapie to sheep. *BMC Vet Res*. 2007;3:20.
- Simmons MM, Moore SJ, Konold T, Thurston L, Terry LA, Thorne L, Lockey R, Vickery C, Hawkins SA, Chaplin MJ, Spiropoulos J. Experimental oral transmission of atypical scrapie to sheep. *Emerg Infect Dis*. 2011;17(5):848-54.
- Simmons MM, Moore SJ, Lockey R, Chaplin MJ, Konold T, Vickery C, Spiropoulos J. Phenotype shift from atypical scrapie to CH1641 following experimental transmission in sheep. *PLoS One*. 2015;10(2):e0117063.
- Smith CB, Booth CJ, Pedersen JA. Fate of prions in soil: a review. *J Environ Qual*. 2011;40(2):449-61.
- Smith JD, Nicholson EM, Greenlee JJ. Evaluation of a combinatorial approach to prion inactivation using an oxidizing agent, SDS, and proteinase K. *BMC Vet Res*. 2013;9:151.
- Smith M, Sherman D. *Goat medicine*. Pennsylvania: Lea and Febiger; 1994. Scrapie; p. 133-5.
- Smith PG, Bradley R. Bovine spongiform encephalopathy (BSE) and its epidemiology. *Br Med Bull*. 2003;66:185-98.
- Sofianidis G, Psychas V, Billinis C, Spyrou V, Argyroudis S, Vlemmas I. Atypical PrPsc distribution in goats naturally affected with scrapie. *J Comp Pathol*. 2008;138(2-3):90-101.
- Sohn HJ, Kim JH, Choi KS, Nah JJ, Joo YS, Jean YH, Ahn SW, Kim OK, Kim DY, Balachandran A. A case of chronic wasting disease in an elk imported to Korea from Canada. *J Vet Med Sci*. 2002;64:855-8.
- Somerville RA, Gentles N. Characterization of the effect of heat on agent strains of the transmissible spongiform encephalopathies. *J Gen Virol*. 2011;92(Pt 7):1738-48.
- Spencer MD, Knight RS, Will RG. First hundred cases of variant Creutzfeldt-Jakob disease: retrospective case note review of early psychiatric and neurological features. *BMJ*. 2002;324:1479-82.
- Spiropoulos J, Hawkins SA, Simmons MM, Bellworthy SJ. Evidence of *in utero* transmission of classical scrapie in sheep. *J Virol*. 2014;88(8):4591-4.
- Spiropoulos J, Lockey R, Sallis RE, Terry LA, Thorne L, Holder TM, Beck KE, Simmons MM. Isolation of prion with BSE properties from farmed goat. *Emerg Infect Dis*. 2011;17(12):2253-61.
- Spraker TR, Gidlewski TL, Balachandran A, VerCauteren KC, Creekmore L, Munger RD. Detection of PrP(CWD) in postmortem rectal lymphoid tissues in Rocky Mountain elk (*Cervus elaphus nelsoni*) infected with chronic wasting disease. *J Vet Diagn Invest*. 2006;18:553-7.
- Spraker TR, VerCauteren KC, Gidlewski T, Schneider DA, Munger R, Balachandran A, O'Rourke KI. Antemortem detection of PrPCWD in preclinical, ranch-raised Rocky Mountain elk (*Cervus elaphus nelsoni*) by biopsy of the rectal mucosa. *J Vet Diagn Invest*. 2009;21(1):15-24.
- Stack M, Jeffrey M, Gubbins S, Grimmer S, González L, Martin S, Chaplin M, Webb P, Simmons M, Spencer Y, Bellerby P, Hope J, Wilesmith J, Matthews D. Monitoring for bovine spongiform encephalopathy in sheep in Great Britain, 1998-2004. *J Gen Virol*. 2006;87:2099-107.
- Suardi S, Vimercati C, Casalone C, Gelmetti D, Corona C, et al. Infectivity in skeletal muscle of cattle with atypical bovine spongiform encephalopathy. *PLoS One*. 2012;7(2):e31449.
- Tamgüney G, Miller MW, Wolfe LL, Sirochman TM, Glidden DV, Palmer C, Lemus A, DeArmond SJ, Prusiner SB. Asymptomatic deer excrete infectious prions in faeces. *Nature*. 2009;461(7263):529-32.
- Tamgüney G, Richt JA, Hamir AN, Greenlee JJ, Miller MW, Wolfe LL, Sirochman TM, Young AJ, Glidden DV, Johnson NL, Giles K, DeArmond SJ, Prusiner SB. Salivary prions in sheep and deer. *Prion*. 2012;6(1):52-61.
- Terry LA, Howells L, Bishop K, Baker CA, Everest S, Thorne L, Maddison BC, Gough KC. Detection of prions in the faeces of sheep naturally infected with classical scrapie. *Vet Res*. 2011;42:65.
- Terry LA, Howells L, Hawthorn J, Edwards JC, Moore SJ, Bellworthy SJ, Simmons H, Lizano S, Estey L, Leathers V, Everest SJ. Detection of PrPsc in blood from sheep infected with the scrapie and bovine spongiform encephalopathy agents. *J Virol*. 2009;83(23):12552-8.
- Terry LA, Jenkins R, Thorne L, Everest SJ, Chaplin MJ, Davis LA, Stack MJ. First case of H-type bovine spongiform encephalopathy identified in Great Britain. *Vet Rec*. 2007;160:873-4.

Transmissible Spongiform Encephalopathies

- The National Creutzfeldt-Jakob Disease Surveillance Unit [CJD Unit], United Kingdom. Variant CJD cases worldwide. CJD Unit, U.K.; 2016 May. Available at: <http://www.cjd.ed.ac.uk/documents/worldfigs.pdf>. Accessed 28 Aug 2016.
- Thomas JG, Chenoweth CE, Sullivan SE. Iatrogenic Creutzfeldt-Jakob disease via surgical instruments. *J Clin Neurosci*. 2013;20(9):1207-12.
- Thomsen BV, Schneider DA, O'Rourke KI, Gidlewski T, McLane J, Allen RW, McIsaac AA, Mitchell GB, Keane DP, Spraker TR, Balachandran A. Diagnostic accuracy of rectal mucosa biopsy testing for chronic wasting disease within white-tailed deer (*Odocoileus virginianus*) herds in North America: effects of age, sex, polymorphism at PRNP codon 96, and disease progression. *J Vet Diagn Invest*. 2012;24(5):878-87.
- Touzeau S, Chase-Topping ME, Matthews L, Lajous D, Eychenne F, Hunter N, Foster JD, Simm G, Elsen JM, Woolhouse ME. Modelling the spread of scrapie in a sheep flock: evidence for increased transmission during lambing seasons. *Arch Virol*. 2006;151:735-51.
- Torres JM, Andréoletti O, Lacroux C, Prieto I, Lorenzo P, Larska M, Baron T, Espinosa JC. Classical bovine spongiform encephalopathy by transmission of H-type prion in homologous prion protein context. *Emerg Infect Dis*. 2011;17(9):1636-44.
- Tyshenko MG. Bovine spongiform encephalopathy and the safety of milk from Canadian dairy cattle. *Vet Rec*. 2007;160:215-8.
- U.K. Food Standards Agency. BSE and beef: new controls explained. Available at: www.food.gov.uk/sites/default/files/multimedia/pdfs/publications/bsebooklet.pdf. Accessed 29 Aug 2016.
- United Kingdom Department for Environment Food and Rural Affairs (DEFRA). Official statistics. Exotic species and domestic cats: TSE surveillance statistics. DEFRA; 2016 Feb. Available at: <https://www.gov.uk/government/statistics/exotic-species-and-domestic-cats-tse-surveillance-statistics>. Accessed 5 Sept 2016.
- U.K. Veterinary Laboratories Agency (VLA). Cattle TSE surveillance statistics. VLA; 5 Apr 2012. Available at: http://vla.defra.gov.uk/science/sci_tse_stats_catt.htm. Accessed 30 Apr 2012.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service [USDA APHIS]. Bovine spongiform encephalopathy (BSE): ongoing surveillance program. Available at: https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/cattle-disease-information/sa_bse/ct_monthly_surv_test/. Accessed 29 Aug 2016.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service [USDA APHIS]. Bovine spongiform encephalopathy (BSE) response plan summary. USDA APHIS; 1998 Oct. Available at: <http://permanent.access.gpo.gov/lps3025/bsesum.pdf>. Accessed 215 Aug 2007.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service [USDA APHIS]. BSE surveillance in the United States. USDA APHIS; 2016 Jun. Available at: https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/cattle-disease-information/sa_bse/ct_surv_in_usa. Accessed 29 Aug 2016.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service [USDA APHIS]. National scrapie eradication program. Available at: https://www.aphis.usda.gov/aphis/ourfocus/animalhealth/animal-disease-information/sheep-and-goat-health/national-scrapie-eradication-program/ct_scrapie_home. Accessed 30 Aug 2016.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service [USDA APHIS]. Scrapie program standards volume 2: scrapie-free flock certification program. USDA APHIS; 2016 May. Available at: https://www.aphis.usda.gov/animal_health/animal_diseases/scrapie/downloads/standards_current.pdf. Accessed 28 Aug. 2016.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service [USDA APHIS]. Centers for Epidemiology and Animal Health. National scrapie surveillance plan. USDA APHIS; 2010 Sept. Available at: https://www.aphis.usda.gov/animal_health/animal_diseases/scrapie/downloads/national_scrapie_surv_plan.pdf. Accessed 30 Aug 2016.
- U.S. Department of Agriculture. Animal and Plant Health Inspection Service [USDA APHIS]. Transmissible mink encephalopathy. USDA APHIS; 2002 Feb. Available at: http://www.aphis.usda.gov/lpa/pubs/fsheet_faq_notice/fs_ahtme.html. Accessed 5 Oct 2008.
- U.S. Department of Agriculture, Animal and Plant Health Inspection Service [USDA APHIS]. Centers for Epidemiology and Animal Health. Highlights of phase II: Scrapie: Ovine slaughter surveillance study 2002-2003 [online]. USDA APHIS; 2004 March. Available at: http://nahms.aphis.usda.gov/sheep/SOSS_highlights.pdf. Accessed 5 Apr 2007.
- U.S. Food and Drug Administration [FDA]. Bovine spongiform encephalopathy. FDA; 2016 Mar. Available at: <http://www.fda.gov/animalveterinary/guidancecomplianceenforcement/complianceenforcement/bovinespongiformencephalopathy/default.htm>. Accessed 30 Aug 2016.
- Vaccari G, Di Bari MA, Morelli L, Nonno R, Chiappini B, Antonucci G, Marcon S, Esposito E, Fazzi P, Palazzini N, Troiano P, Petrella A, Di Guardo G, Agrimi U. Identification of an allelic variant of the goat PrP gene associated with resistance to scrapie. *J Gen Virol*. 2006;87:1395-402.
- Vaccari G, Panagiotidis CH, Acin C, Peletto S, Barillet F, et al. State-of-the-art review of goat TSE in the European Union, with special emphasis on PRNP genetics and epidemiology. *Vet Res*. 2009;40(5):48.
- Vascellari M, Nonno R, Mutinelli F, Bigolaro M, Di Bari MA, Melchioti E, Marcon S, D'Agostino C, Vaccari G, Conte M, De Grossi L, Rosone F, Giordani F, Agrimi U. PrPSc in salivary glands of scrapie-affected sheep. *J Virol*. 2007;81:4872-6.
- VerCauteren KC, Pilon JL, Nash PB, Phillips GE, Fischer JW. Prion remains infectious after passage through digestive system of American crows (*Corvus brachyrhynchos*). *PLoS One*. 2012;7(10):e45774.
- Williams ES. Chronic wasting disease. *Vet Pathol*. 2005;42:530-49.
- Williams ES, Miller MW. Transmissible spongiform encephalopathies in non-domestic animals: origin, transmission and risk factors. *Rev Sci Tech*. 2003;22:145-56.

Transmissible Spongiform Encephalopathies

- World Organization for Animal Health [OIE]. Manual of diagnostic tests and vaccines [online]. Paris: OIE; 2016. Bovine spongiform encephalopathy. Available at: http://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/2.04.05_BSE.pdf. Accessed 26 Aug 2016.
- World Organization for Animal Health [OIE]. Manual of diagnostic tests and vaccines [online]. Paris: OIE; 2016. Scrapie. Available at: http://www.oie.int/fileadmin/Home/eng/Health_standards/tahm/2.07.12_SCRAPIE.pdf. Accessed 20 Aug 2016.
- World Organization for Animal Health [OIE]. Number of reported cases of bovine spongiform encephalopathy (BSE) in farmed cattle worldwide (excluding the United Kingdom). Available at: <http://www.oie.int/animal-health-in-the-world/bse-specific-data/number-of-reported-cases-worldwide-excluding-the-united-kingdom/>. Accessed 29 Aug 2016.
- World Organization for Animal Health [OIE]. Number of cases of bovine spongiform encephalopathy (BSE) reported in the United Kingdom. Available at: <http://www.oie.int/animal-health-in-the-world/bse-specific-data/number-of-cases-in-the-united-kingdom/>. Accessed 29 Aug 2016.
- World Organization for Animal Health [OIE]. Terrestrial animal health code [online]. Paris: OIE; 2016. Bovine spongiform encephalopathy. Available at: http://www.oie.int/index.php?id=169&L=0&htmfile=chapitre_bse.htm. Accessed 29 Aug 2016.
- World Organization for Animal Health [OIE]. World animal health information database (WAHIS) interface. Scrapie. OIE; 2015. Available at: http://www.oie.int/wahis_2/public/wahid.php/Wahidhome/Home. Accessed 28 Aug 2016.
- Wyatt JM, Pearson GR, Gruffydd-Jones TJ. Feline spongiform encephalopathy. *Feline Pract.* 1993;21:7-9.
- Xu S, Reuter T, Gilroyed BH, Mitchell GB, Price LM, Dudas S, Braithwaite SL, Graham C, Czub S, Leonard JJ, Balachandran A, Neumann NF, Belosevic M, McAllister TA. Biodegradation of prions in compost. *Environ Sci Technol.* 2014;48(12):6909-18.
- Yokoyama T, Okada H, Murayama Y, Masujin K, Iwamaru Y, Mohri S. Examination of the offspring of a Japanese cow affected with L-type bovine spongiform encephalopathy. *J Vet Med Sci.* 2011;73(1):121-3.
- Young S, Slocombe RF. Prion-associated spongiform encephalopathy in an imported Asiatic golden cat (*Catopuma temminckii*). *Aust Vet J.* 2003;81:295-6.
- Yuan Q, Eckland T, Telling G, Bartz J, Bartelt-Hunt S. Mitigation of prion infectivity and conversion capacity by a simulated natural process--repeated cycles of drying and wetting. *PLoS Pathog.* 2015;11(2):e1004638.
- Zanusso G, Casalone C, Acutis P, Bozzetta E, Farinazzo A, Gelati M, Fiorini M, Forloni G, Sy MS, Monaco S, Caramelli M. Molecular analysis of iatrogenic scrapie in Italy. *J Gen Virol.* 2003;84:1047-52.
- Zanusso G, Nardelli E, Rosati A, Fabrizi G, Ferrari S, Carteri A, De Simone F, Rizzuto N, Monaco S. Simultaneous occurrence of spongiform encephalopathy in a man and his cat in Italy. *Lancet.* 1998;352:1116-7.

*Link defunct